

Initial-boundary value problem for distributed order time-fractional diffusion equations

Zhiyuan LI*, Yavar KIAN† and Éric SOCCORSI‡

Abstract

We examine initial-boundary value problems for diffusion equations with distributed order time-fractional derivatives. We prove existence and uniqueness results for the weak solution to these systems, together with its continuous dependency on initial value and source term. Moreover, we provide energy estimates of the solution for small and large times. Finally, under suitable assumption on the source term, we establish that the solution is analytic in time.

1 Introduction

The time-fractional diffusion model of constant order (CO) α , $\partial_t^\alpha u - \Delta u = f$, has received great attention within the last few decades from numerous applied scientists, see e.g. [2, 3, 7, 14], due to its relevance for modeling anomalous diffusion processes whose mean square displacement (MSD) scales like t^α as the time variable t goes to infinity. But, more recently, it was noticed in [4, 21, 26] and the references therein that several application areas such as polymer physics or kinetics of particles moving in quenched random force fields, exhibit ultraslow diffusion phenomena whose MSD is of logarithmic growth. There are several approaches for modeling such processes. One of them uses time-fractional diffusion equations of distributed order (DO), see e.g. [12, 19, 24], but we also mention that a diffusion model with variable fractional order time-derivative was proposed in [27] to depict ultraslow diffusion processes. In this paper, we are concerned with the DO fractional diffusion model. More precisely, we consider the following initial-boundary value problem (IBVP),

$$\begin{cases} \mathbb{D}_t^{(\mu)} u + \mathcal{A}u = F & \text{in } Q := \Omega \times (0, T), \\ u(0, \cdot) = u_0 & \text{in } \Omega, \\ u = 0 & \text{on } \Sigma := \partial\Omega \times (0, T), \end{cases} \quad (1)$$

where Ω is an open bounded domain in \mathbb{R}^d , $d \geq 2$, with $\mathcal{C}^{1,1}$ boundary $\partial\Omega$, and \mathcal{A} is the following symmetric and uniformly elliptic operator

$$\mathcal{A}\varphi(x) := - \sum_{i,j=1}^d \partial_{x_i} (a_{ij}(x) \partial_{x_j} \varphi(x)) + q(x) \varphi(x),$$

Manuscript last updated: March 14, 2019.

† Aix-Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France.

E-mails : yavar.kian@univ-amu.fr (Yavar Kian) and eric.soccorsi@univ-amu.fr (Éric Soccorsi).

* Graduate School of Mathematical Sciences, the University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo 153-8914, Japan.

E-mail: zyli@ms.u-tokyo.ac.jp.

Funding: The work of ZL was partially supported by Grant-in-Aid for Scientific Research (S) 15H05740, JSPS (ZL). The work of YK and ÉS was partially supported by the Agence Nationale de la Recherche under grant ANR-17-CE40-0029.

associated with a suitable real-valued electric potential q and symmetric coefficients $a_{ij} = a_{ji}$, $1 \leq i, j \leq d$, fulfilling the uniform ellipticity condition

$$\sum_{i,j=1}^d a_{ij}(x)\xi_i\xi_j \geq c_a|\xi|^2, \quad x \in \bar{\Omega}, \quad \xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d, \quad (2)$$

for some positive constant c_a . Here, $\mathbb{D}_t^{(\mu)}$ denotes the distributed order fractional derivative

$$\mathbb{D}_t^{(\mu)}\varphi(t) := \int_0^1 \partial_t^\alpha \varphi(t) \mu(\alpha) d\alpha,$$

induced by the non-negative weight function $\mu \in L^\infty(0,1)$, and ∂_t^α is the Caputo derivative of order α , defined by:

$$\partial_t^\alpha \varphi(t) := \begin{cases} \varphi(t) & \text{if } \alpha = 0, \\ \frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{\varphi(\tau)}{(t-\tau)^\alpha} d\tau & \text{if } \alpha \in (0,1), \\ \varphi'(t) & \text{if } \alpha = 1, \end{cases}$$

the symbol Γ denoting the usual Gamma function.

From a mathematical viewpoint, the forward problem associated with these equations was investigated in [12, 17, 18, 20]. Namely, the fundamental solution to the Cauchy problem for both ordinary and partial distributed order fractional differential equations with continuous weight function was derived and investigated in detail in [12]. A uniqueness result for the solution to diffusion equations of DO was derived in [18] with the aid of an appropriate maximum principle and a formal solution was constructed by means of the Fourier method of variables separation. Unfortunately, there is no proof available in [18] of the convergence of the series describing this formal solution. Further, explicit strong solutions (and stochastic analogues) to DO time-fractional diffusion equations with Dirichlet boundary conditions were built in [20] for C^1 -weight functions. For the asymptotic behavior we refer to [17], where logarithmic decay of the solution to DO diffusion equations was established for $t \rightarrow \infty$, while this solution scales at best like $(t|\log t|)^{-1}$ as $t \rightarrow 0$. These results are in sharp contrast with the ones derived for single- and multi-term time-fractional diffusion equations, see e.g. [15, 25], which seem unable to capture the time asymptotic trends of ultraslow diffusion processes.

Formally, single- or multi-term time-fractional diffusion equations can be seen as DO time-fractional diffusion equations associated with a density function of the form $\mu = \sum_{j=1}^\ell q_j \delta(\cdot - \alpha_j)$, where δ is the Dirac-delta function. The definition of a weak solution for single- or multi-term time-fractional diffusion equations was recently introduced in [6, 8] by taking advantage of the fact that the domain of the $L^2(0,T)$ -realization of the operator ∂_t^α , $\alpha \in (0,1)$, is embedded in the fractional Sobolev space of order α , $H^\alpha(0,T)$. However, as it is still unclear whether the domain of $\mathbb{D}_t^{(\mu)}$ can be described by fractional Sobolev spaces, the scheme developed in [6, 8] does not seem to be relevant for DO time-fractional diffusion equations. Another approach, initiated by [28] and recently applied to DO time-fractional diffusion equations in [13], is to consider (1) as an abstract evolutionary integro-differential equation. This strategy is suitable for both autonomous and non-autonomous equations but there is a serious inconvenience to this method, arising from the dependency of the functional space of the solution on the kernel function of the corresponding integro-differential operator, which causes numerous technical difficulties in performing computations based on this model.

In this paper, since the system under study is autonomous (the coefficients appearing in (1) are all space-dependent only), we rather follow the idea of [10, 11] and characterize the weak solution to (1) as the original of the solution to the Laplace transform of (1) with respect to the time variable. With reference to the analysis carried out in [15] for multi term CO

time-fractional diffusion equations, we aim to study the existence, uniqueness and regularity properties, and the stability with respect to the diffusion coefficients and the weight function μ , of a weak solution to (1).

1.1 Settings

In this paper, we assume that $q \in L^\kappa(\Omega)$ for some $\kappa \in (d, +\infty)$, is non-negative, i.e.

$$q(x) \geq 0, \quad x \in \Omega, \quad (3)$$

and that the coefficients $a_{i,j} = a_{j,i} \in C^1(\overline{\Omega}, \mathbb{R})$, $1 \leq i, j \leq d$, satisfy the ellipticity condition (2).

We denote by A the operator generated in $L^2(\Omega)$ by the quadratic form

$$u \mapsto \sum_{i,j=1}^d \int_{\Omega} a_{i,j}(x) \partial_{x_i} u(x) \partial_{x_j} \overline{u(x)} dx + \int_{\Omega} q(x) |u(x)|^2 dx, \quad u \in H_0^1(\Omega).$$

Due to (2), A is selfadjoint in $L^2(\Omega)$ and acts as the operator \mathcal{A} on its domain $D(A) := H_0^1(\Omega) \cap H^2(\Omega)$, in virtue of [10, Section 2.1].

Moreover, since $H_0^1(\Omega)$ is compactly embedded in $L^2(\Omega)$, the resolvent of A is compact in $L^2(\Omega)$, hence the spectrum of A is purely discrete. We denote by $\{\lambda_n, n \in \mathbb{N}\}$, where $\mathbb{N} := \{1, 2, \dots\}$, the sequence of the eigenvalues of A arranged in non-decreasing order and repeated according to their multiplicity. In light of (2)-(3), we have

$$\lambda_n \geq c_a > 0, \quad n \in \mathbb{N}. \quad (4)$$

For further use, we introduce an orthonormal basis $\{\varphi_n, n \in \mathbb{N}\}$ of eigenfunctions of A in $L^2(\Omega)$, such that

$$A\varphi_n = \lambda_n \varphi_n, \quad \varphi_n \in D(A), \quad n \in \mathbb{N}.$$

1.2 Weak solution

As already mentioned in the introduction, the usual definition given in [6] of a weak solution to CO time-fractional diffusion equations, is not suitable for DO time-fractional diffusion equations. Hence we rather follow the strategy implemented in [10] (which is by means of the Laplace transform of tempered distributions), that is recalled below.

Let $\mathcal{S}'(\mathbb{R}, L^2(\Omega)) := \mathcal{B}(\mathcal{S}(\mathbb{R}, L^2(\Omega)), \mathbb{R})$ be the space dual to $\mathcal{S}(\mathbb{R}; L^2(\Omega))$, put $\mathbb{R}_+ := [0, +\infty)$, and denote by $\mathcal{S}'(\mathbb{R}_+, L^2(\Omega)) := \{v \in \mathcal{S}'(\mathbb{R}, L^2(\Omega)); \text{supp } v \subset \mathbb{R}_+ \times \overline{\Omega}\}$ the set of distributions in $\mathcal{S}'(\mathbb{R}, L^2(\Omega))$ that are supported in $\mathbb{R}_+ \times \overline{\Omega}$. Otherwise stated, $v \in \mathcal{S}'(\mathbb{R}, L^2(\Omega))$ lies in $\mathcal{S}'(\mathbb{R}_+, L^2(\Omega))$ if and only if $\langle v, \varphi \rangle_{\mathcal{S}'(\mathbb{R}, L^2(\Omega)), \mathcal{S}(\mathbb{R}, L^2(\Omega))} = 0$ whenever $\varphi \in \mathcal{S}(\mathbb{R}, L^2(\Omega))$ vanishes in $\mathbb{R}_+ \times \overline{\Omega}$. As a consequence we have

$$\langle v(\cdot, x), \varphi \rangle_{\mathcal{S}'(\mathbb{R}), \mathcal{S}(\mathbb{R})} = \langle v(\cdot, x), \psi \rangle_{\mathcal{S}'(\mathbb{R}), \mathcal{S}(\mathbb{R})}, \quad \varphi, \psi \in \mathcal{S}(\mathbb{R}),$$

provided $\varphi = \psi$ in \mathbb{R}_+ . Further, we say that $\varphi \in \mathcal{S}(\mathbb{R}_+)$ if φ is the restriction to \mathbb{R}_+ of a function $\tilde{\varphi} \in \mathcal{S}(\mathbb{R})$. Then, for a.e. $x \in \Omega$, we set

$$\langle v(\cdot, x), \varphi \rangle_{\mathcal{S}'(\mathbb{R}_+), \mathcal{S}(\mathbb{R}_+)} := x \mapsto \langle v(\cdot, x), \tilde{\varphi} \rangle_{\mathcal{S}'(\mathbb{R}), \mathcal{S}(\mathbb{R})}, \quad v \in \mathcal{S}'(\mathbb{R}_+, L^2(\Omega)).$$

Notice that $\tilde{\varphi}$ may be any function in $\mathcal{S}(\mathbb{R})$ such that $\tilde{\varphi}(t) = \varphi(t)$ for all $t \in \mathbb{R}_+$.

For all $p \in \mathbb{C}_+ := \{z \in \mathbb{C}; \text{Re } z \in (0, +\infty)\}$, where $\text{Re } z$ denotes the real part of z , we put

$$e_p(t) := \exp(-pt), \quad t \in \mathbb{R}_+.$$

Evidently, e_p lies in $\mathcal{S}(\mathbb{R}_+)$ so we can define the Laplace transform $\mathcal{L}[v]$ of $v \in \mathcal{S}'(\mathbb{R}_+, L^2(\Omega))$, with respect to t , as the family of mappings

$$\mathcal{L}[v](p) : x \in \Omega \mapsto \langle v(\cdot, x), e_p \rangle_{\mathcal{S}'(\mathbb{R}_+), \mathcal{S}(\mathbb{R}_+)}, \quad p \in \mathbb{C}_+.$$

Notice that $p \mapsto \mathcal{L}[v](p) \in \mathcal{C}^\infty(\mathbb{C}_+, L^2(\Omega))$.

Inspired by [10, Definition 2.2] we may now introduce the weak solution to (1) as follows.

Definition 1.1. *Let $\mu \in L^\infty(0, 1; \mathbb{R}_+)$, let $u_0 \in L^2(\Omega)$ and, depending on whether $T \in (0, +\infty)$ or $T = +\infty$, let $F \in L^\infty(0, T; L^2(\Omega))$ or let $t \mapsto (1+t)^{-m}F(t, \cdot) \in L^\infty(\mathbb{R}_+, L^2(\Omega))$ for some $m \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$. We say that u is a weak solution to the IBVP (1) if u is the restriction to Q of a distribution $v \in \mathcal{S}'(\mathbb{R}_+, L^2(\Omega))$, i.e. $u = v|_Q$, whose Laplace transform $\widehat{v} := \mathcal{L}[v]$ fulfills*

$$(A + sw(s))\widehat{v}(s) = w(s)u_0 + \widehat{F}(s), \quad s \in \mathbb{C}_+, \quad (5)$$

with $w(s) := \int_0^1 s^{\alpha-1} \mu(\alpha) d\alpha$ and $\widehat{F}(s) := \mathcal{L}[F \mathbb{1}_{(0, T)}](s) = \int_0^T e^{-st} F(t, \cdot) dt$. Here $\mathbb{1}_{(0, T)}$ stands for the characteristic function of $(0, T)$.

Notice that Equation (5) yields $\widehat{v}(s) \in D(A)$ for all $s \in \mathbb{C}_+$.

Remark 1.1. *A solution to the IBVP (1), as defined by Definition 1.1, is necessarily unique. Indeed, if u_1 and u_2 are two such solutions then there are two tempered distributions v_1 and v_2 in $\mathcal{S}'(\mathbb{R}_+, L^2(\Omega))$ such that $u_i = (v_i)|_Q$ for $i = 1, 2$, $(\widehat{v}_1 - \widehat{v}_2)(s) \in H_0^1(\Omega)$ and*

$$(A + sw(s))(\widehat{v}_1 - \widehat{v}_2)(s) = 0, \quad s \in \mathbb{C}_+.$$

Moreover, since A is a positive operator, $-sw(s)$ is in the resolvent set of A for each $s \in \mathbb{C}_+$. Therefore the above identity yields

$$(\widehat{v}_1 - \widehat{v}_2)(s) = 0, \quad s \in \mathbb{C}_+,$$

so we get $v_1 = v_2$ by injectivity of the Laplace transform, and consequently $u_1 = u_2$.

In the coming section we state several existence and uniqueness results for the weak solution to (1).

1.3 Main results

We first address the case where the initial state $u_0 \in L^2(\Omega)$. The corresponding result is as follows.

Theorem 1.1. *Let $\mu \in L^\infty(0, 1)$ be non-negative and fulfill the following condition:*

$$\exists \alpha_0 \in (0, 1), \exists \delta \in (0, \alpha_0), \forall \alpha \in (\alpha_0 - \delta, \alpha_0), \mu(\alpha) \geq \frac{\mu(\alpha_0)}{2} > 0. \quad (6)$$

Depending on whether $T \in (0, +\infty)$ or $T = +\infty$, we assume either that $F \in L^\infty(0, T; L^2(\Omega))$ or that $t \mapsto (1+t)^{-m}F(t, \cdot) \in L^\infty(\mathbb{R}_+, L^2(\Omega))$ for some natural number $m \in \mathbb{N}_0$.

Then, for all $u_0 \in L^2(\Omega)$, there exists a unique weak solution $u \in \mathcal{C}((0, T], L^2(\Omega))$ to (1). Moreover, we have $u \in \mathcal{C}([0, T], L^2(\Omega))$ provided $u_0 = 0$.

Here and in the remaining part of this text, the notation $(0, T]$ (resp., $[0, T]$) stands for $(0, +\infty)$ (resp., $[0, +\infty)$) in the particular case where $T = +\infty$. For a finite final time T , we have the following improved regularity result.

Theorem 1.2. *Fix $T \in (0, +\infty)$ and let μ be the same as in Theorem 1.1.*

(a) Pick $u_0 \in D(A^\gamma)$ for $\gamma \in (0, 1]$, and let $F = 0$. Then, the unique weak solution u to (1) lies in $\mathcal{C}([0, T], L^2(\Omega)) \cap \mathcal{C}((0, T], H_0^1(\Omega) \cap H^2(\Omega)) \cap \mathcal{C}^1((0, T], L^2(\Omega))$ and satisfies the two following estimates

$$\|u(t)\|_{H^2(\Omega)} \leq Ce^T \|u_0\|_{D(A^\gamma)} t^{\gamma-1}, \quad t \in (0, T], \quad (7)$$

and

$$\|\partial_t u(t)\|_{L^2(\Omega)} \leq Ce^T \|u_0\|_{D(A^\gamma)} t^{-\beta}, \quad \beta \in (1 - \alpha_0 \gamma, 1), \quad t \in (0, T], \quad (8)$$

where C is a positive constant which is independent of T , t and u_0 .

(b) Assume that $u_0 = 0$ and $F \in L^\infty(0, T; L^2(\Omega))$. Then, for all $\kappa \in [0, 1]$ and all $p \in \left[1, \frac{1}{1 - \alpha_0(1 - \kappa)}\right)$, the unique weak solution u to (1) lies in $L^p(0, T; H^{2\kappa}(\Omega))$. Moreover, there exists a positive constant C , independent of T and F , such that we have

$$\|u\|_{L^p(0, T; H^{2\kappa}(\Omega))} \leq Ce^{CT} \|F\|_{L^1(0, T; L^2(\Omega))}. \quad (9)$$

Estimate (8) holds for $\gamma \in (0, 1]$ and $\alpha_0 \in (0, 1)$, hence for $1 - \alpha_0 \gamma \in (0, 1)$. Thus the right-hand side of the inequality scales at best as $t^{-(1 - \alpha_0 \gamma)}$, which is coherent with the analysis carried out in [22, 25], indicating that the $L^2(\Omega)$ -norm of the first derivative of the solution to fractional diffusion equations with CO $\alpha \in (0, 1)$, initial state in $D(A^\gamma)$ and identically zero source term, may blow up like $t^{-1 + \alpha \gamma}$ as t approaches 0. Such a singular behavior at initial time is compatible with [1, Theorem 7.3], where the weak solution to time-fractional PDEs is defined in a variational sense.

Moreover, we notice that the second claim of Theorem 1.2 states that $u \in L^p(0, T; D(A^\kappa))$ for suitable values of p , provided $u_0 = 0$ and $F \in L^\infty(0, T; L^2(\Omega))$. Here κ can be arbitrarily close to 1 without actually becoming 1, except for the special case treated by the following result, where the density function μ vanishes in a neighborhood of the endpoint of the interval $(0, 1)$.

Theorem 1.3. *Let $\mu \in \mathcal{C}([0, 1], \mathbb{R}_+)$ satisfy (6). Assume moreover that there exists $\alpha_1 \in (\alpha_0, 1)$, where α_0 is the same as in (6), such that*

$$\mu(\alpha) = 0, \quad \alpha \in (\alpha_1, 1). \quad (10)$$

Then, for $u_0 = 0$ and $F \in L^\infty(0, T; L^2(\Omega))$, there exists a unique solution $u \in L^2(0, T; H^2(\Omega))$ to the IBVP (1), satisfying

$$\|u\|_{L^2(0, T; H^2(\Omega))} \leq C \|F\|_{L^2(Q)},$$

for some positive constant C , which is independent of F .

The following result claims that the solution to (1) is analytic in time, provided the source term can be holomorphically extended to a neighborhood of the positive real axis. This statement is of great interest in the analysis of inverse coefficient problems associated with time-fractional diffusion equations, see e.g. [9, 10, 16].

Theorem 1.4. *Let the conditions of Theorem 1.1 be satisfied with $T = +\infty$. Assume moreover that there exists $\rho \in (0, +\infty)$ such that the source term $t \mapsto F(t, \cdot)$ is extendable to a holomorphic function of the half-strip $S_\rho := \{z \in \mathbb{C}; \operatorname{Re} z \in (-\rho, +\infty), \operatorname{Im} z \in (-\rho, \rho)\}$ into $L^2(\Omega)$, where we recall that $\operatorname{Re} z$ (resp., $\operatorname{Im} z$) stands for the real (resp., imaginary) part of z . Then, the weak-solution $t \mapsto u(t, \cdot)$ to (1), given by Theorem 1.1, can be extended to an analytic function of $(0, +\infty)$ into $L^2(\Omega)$.*

The last result of this paper is similar to [15, Theorem 2.3], which was established in the framework of multi-terms time-fractional diffusion equations with positive constant coefficients.

It is useful for the optimization approach to the inverse problem of determining the weight function μ together with the diffusion matrix $a := (a_{i,j})_{1 \leq i,j \leq d}$ and the electric potential q , appearing in the definition of the operator \mathcal{A} , by extra data of the solution to (1). Namely, we claim for all *a priori* fixed $M \in (0, +\infty)$ that the weak solution to the IBVP (1) associated with $F = 0$, depends Lipschitz continuously on (μ, a, q) in $\mathcal{W}(M) \times \mathcal{D}(M) \times \mathcal{Q}(M)$, where

$$\mathcal{W}(M) := \left\{ \mu \in L^\infty(0, 1; \mathbb{R}_+) \text{ satisfying (6) and } \|\mu\|_{L^\infty(0,1)} \leq M \right\}$$

is the set of admissible weight functions,

$$\mathcal{D}(M) := \{a = a^T = (a_{i,j})_{1 \leq i,j \leq d} \in C^1(\bar{\Omega}, \mathbb{R}^{d^2}) \text{ fulfilling (2) and } \|a\|_{C^1(\bar{\Omega})} \leq M\},$$

with a^T the transpose matrix of a , denotes the set of admissible diffusion matrices, and

$$\mathcal{Q}(M) := \{q \in C^0(\bar{\Omega}) \text{ obeying (3) and } \|q\|_{C^0(\bar{\Omega})} \leq M\},$$

is the set of admissible electric potentials.

Theorem 1.5. *Let $T \in (1, +\infty)$, fix $M \in (0, +\infty)$ and pick (μ, a, q) and $(\tilde{\mu}, \tilde{a}, \tilde{q})$ in $\mathcal{W}(M) \times \mathcal{D}(M) \times \mathcal{Q}(M)$. For $u_0 \in D(A)$ such that $\|u_0\|_{D(A)} \leq M$, let u (resp., \tilde{u}) denote the weak solution to the IBVP (1) (resp., the IBVP (1) where $(\tilde{\mu}, \tilde{a}, \tilde{q})$ is substituted for (μ, a, q)) with identically zero source term, given by Statement (a) in Theorem 1.2. Assume moreover that μ and $\tilde{\mu}$ coincide in a neighborhood of 1, i.e. that there exists $\alpha_1 \in (\alpha_0, 1)$ such that*

$$\mu(\alpha) = \tilde{\mu}(\alpha), \quad \alpha \in (\alpha_1, 1). \quad (11)$$

Then we have $u - \tilde{u} \in L^p(0, T; H^{2\kappa}(\Omega))$ for all $\kappa \in (0, 1)$ and all $p \in \left[1, \frac{1}{1-\alpha_0(1-\kappa)}\right)$, and the estimate

$$\|u - \tilde{u}\|_{L^p(0,T;H^{2\kappa}(\Omega))} \leq C \left(\|\mu - \tilde{\mu}\|_{L^\infty(0,1)} + \|a - \tilde{a}\|_{C^1(\bar{\Omega})} + \|q - \tilde{q}\|_{C^0(\bar{\Omega})} \right), \quad (12)$$

for some constant $C \in (0, +\infty)$ depending only on $T, M, \alpha_0, \delta, \alpha_1, \kappa, p$ and c_a .

1.4 Generalization of the results

In Theorems 1.1, 1.2 and 1.3, the function F is assumed to be bounded in time over $(0, T)$ but it turns out that in many applications the relevant source term appearing in (1) lies in $L^1(0, T; L^2(\Omega))$. In such a case the Definition 1.1 of a weak solution to the IBVP (1) is no longer valid but it can be easily adapted to the framework of $F \in L^1(0, T; L^2(\Omega))$ with a density argument we make precise below.

Let us start by introducing the following notation: for all $\varphi \in L^\infty(0, T; L^2(\Omega))$ we denote by $u(\varphi)$ the $L^1(0, T; H_0^1(\Omega))$ -solution to the system (1) associated with $u_0 = 0$ and $F = \varphi$, which is given by Statement (b) in Theorem 1.2 with $p = 1$ and $\kappa = \frac{1}{2}$. Then, for $T \in (0, +\infty)$, $u_0 = 0$ and $F \in L^1(0, T; L^2(\Omega))$, we say that $u \in L^1(0, T; H_0^1(\Omega))$ is a weak solution to (1) if u is the $L^1(0, T; H^1(\Omega))$ -limit of $\{u(F_n), n \in \mathbb{N}\}$, where $\{F_n, n \in \mathbb{N}\} \in L^\infty(0, T; L^2(\Omega))^{\mathbb{N}}$ is an approximation sequence of F in $L^1(0, T; L^2(\Omega))$, i.e.

$$\lim_{n \rightarrow \infty} \|u - u(F_n)\|_{L^1(0,T;H^1(\Omega))} = 0, \quad (13)$$

where $F_n \in L^\infty(0, T; L^2(\Omega))$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} \|F - F_n\|_{L^1(0,T;L^2(\Omega))} = 0$.

Notice that the above definition is meaningful in the sense that, firstly, there exists a unique function $u \in L^1(0, T; H_0^1(\Omega))$ obeying (13), and secondly, u depends only on F and not on the choice of the converging sequence $\{F_n, n \in \mathbb{N}\}$. Indeed, the first claim follows from

the fact that $\{u(F_n), n \in \mathbb{N}\}$ is a Cauchy sequence in $L^1(0, T; H_0^1(\Omega))$. This can be seen from the estimate $\|u(F_{n+k}) - u(F_n)\|_{L^1(0, T; H^1(\Omega))} \leq Ce^{CT} \|u(F_{n+k}) - u(F_n)\|_{L^1(0, T; L^2(\Omega))}$ arising for all n and k in \mathbb{N} by applying (9) with $p = 1$ and $\kappa = \frac{1}{2}$ to the $L^1(0, T; H_0^1(\Omega))$ -solution $u(F_{n+k}) - u(F_n)$ of the system (1) associated with $u_0 = 0$ and $F = F_{n+k} - F_n$. Moreover, if $\{\tilde{F}_n, n \in \mathbb{N}\}$ is another $L^\infty(0, T; L^2(\Omega))$ -sequence fulfilling $\lim_{n \rightarrow \infty} \tilde{F}_n = F$ in $L^1(0, T; L^2(\Omega))$, then we have $\|u(F_n) - u(\tilde{F}_n)\|_{L^1(0, T; H^1(\Omega))} \leq Ce^{CT} \|F_n - \tilde{F}_n\|_{L^1(0, T; L^2(\Omega))}$ from (9) with $p = 1$ and $\kappa = \frac{1}{2}$, since $u(F_n) - u(\tilde{F}_n)$ is a solution to the IBVP (1) associated with $u_0 = 0$ and $F = F_n - \tilde{F}_n$. As a consequence Theorem 1.1 with $T \in (0, +\infty)$ and Theorem 1.2 remain valid with $F \in L^1(0, T; L^2(\Omega))$. In particular, we infer from (9) that the solution u to (1) associated with $u_0 = 0$ and $F \in L^1(0, T; L^2(\Omega))$, fulfills

$$\|u\|_{L^p(0, T; H^{2\kappa}(\Omega))} \leq Ce^{CT} \|F\|_{L^1(0, T; L^2(\Omega))}, \quad \kappa \in [0, 1), \quad p \in \left[1, \frac{1}{1 - \alpha_0(1 - \kappa)}\right). \quad (14)$$

Similarly, it is apparent that the statement of Theorem 1.3 still holds for $T \in (0, +\infty)$ and $F \in L^2(Q)$.

1.5 Brief comments and outline

To our best knowledge, the only mathematical paper besides this one, dealing with the existence and uniqueness issues for solutions to DO fractional diffusion equations, is [13]. But the analysis carried out in [13] is different from the one of the present paper in many aspects. As already mentioned in Section 1, the approach of [13] is variational whereas we study the original function of the solution to the Laplace transform of (1). This allows us to show existence of a unique weak solution to (1) within the class $\mathcal{C}((0, T], L^2(\Omega))$ (and even $\mathcal{C}([0, T], L^2(\Omega))$ if $u_0 = 0$), whereas the solution exhibited in [13] lies in $L^2(Q)$. Similarly, it is unclear whether the improved regularity estimates (7)-(9) or the time analyticity of the weak solution can be derived from the scheme of [13]. This being said, we point out that it is still unclear whether the solution defined in this text and the variational solution of [13] coincide in the case where both notions can be applied, and we stress once more that the approach of [13] applies to non-autonomous systems, which is not the case of the analysis presented in this article.

Finally, we point out that Definition 1.1 of a weak solution to (1) (as the original function of the solution to the Laplace transform with respect to the time variable of this system) is inspired by the analysis carried out in [10], which is concerned with space dependent variable order (VO) time-fractional diffusion equations. It is well known that the weak solution to CO time-fractional diffusion equations can be expressed in terms of Mittag-Leffler functions, see e.g. [5, 11]. Nevertheless, such an explicit representation formula is no longer valid for DO or space-dependent VO time-fractional diffusion equations, as the inversion method of the Laplace transform is technically more involved in these two cases. This specific difficulty arising from the non-constancy of the order of DO or VO time-fractional equations is the main difference with the analysis of their CO counterpart.

The article is organized as follows. In Section 2 we prove Theorem 1.1 by showing existence of a unique weak solution to the IBVP (1), enjoying a Duhamel-like representation formula. Section 3 contains the proof of Theorems 1.2 and 1.3, based on careful analysis of the above mentioned representation of the solution. In Section 4 we establish the time analytic property of the solution to (1), claimed in Theorem 1.4. Next, in Section 5, we prove Theorem 1.5 stating that the weak solution to (1) depends continuously on the distributed order weight function, the diffusion coefficients and the electric potential. Finally, in the appendix presented in Section 6, we collect the proof of an auxiliary result used in the derivation of Theorem 1.3.

2 Representation of the solution: proof of Theorem 1.1

The proof of Theorem 1.1 is based on an effective representation of the solution to the IBVP (1), that is derived in Section 2.2 and presented in Proposition 2.1. As a preamble, several useful properties of the function $s \mapsto w(s) = \int_0^1 s^{\alpha-1} \mu(\alpha) d\alpha$, that are needed by the proofs of Proposition 2.1 and Theorems 1.1 and 1.3, are collected in the coming section.

2.1 Three auxiliary results on w

We start by lower bounding $|sw(s) + \lambda|$ with respect to λ , uniformly in $s \in \mathbb{C} \setminus \mathbb{R}_-$ and $\lambda \in (0, +\infty)$, where $\mathbb{R}_- := (-\infty, 0]$.

Lemma 2.1. *Let $\mu \in L^\infty(0, 1)$ be non-negative. Then, for all $\lambda \in (0, +\infty)$ and all $s = re^{\pm i\beta}$, where $r \in (0, +\infty)$ and $\beta \in [0, \pi)$, we have*

$$|sw(s) + \lambda| \geq C_\beta \lambda \text{ with } C_\beta := \begin{cases} 1 & \text{if } \beta \in [0, \frac{\pi}{2}], \\ \frac{\sin \beta}{2} & \text{if } \beta \in (\frac{\pi}{2}, \pi). \end{cases} \quad (15)$$

Moreover, if $\beta \in (\frac{\pi}{2}, \pi)$, then we have in addition:

$$\frac{\lambda^\nu |sw(s)|^{1-\nu}}{|sw(s) + \lambda|} \leq \frac{2}{\sin \beta}, \quad \nu \in [0, 1]. \quad (16)$$

Proof. We start with (15). The case of $\beta \in [0, \frac{\pi}{2}]$ is easily treated, as we have

$$\operatorname{Re}(sw(s)) = \int_0^1 r^\alpha \cos(\alpha\beta) \mu(\alpha) d\alpha \geq 0,$$

since μ is non-negative, and consequently $\operatorname{Re}(sw(s) + \lambda) \geq \lambda$. In order to examine the case where $\beta \in (\frac{\pi}{2}, \pi)$, we put

$$f(s) := \int_0^{\frac{\pi}{2\beta}} r^\alpha \cos(\alpha\beta) \mu(\alpha) d\alpha \text{ and } g(s) := \int_{\frac{\pi}{2\beta}}^1 r^\alpha |\cos(\alpha\beta)| \mu(\alpha) d\alpha, \quad (17)$$

in such a way that we have

$$\operatorname{Re}(sw(s)) = f(s) - g(s), \quad (18)$$

with

$$f(s) \geq 0 \text{ and } 0 \leq g(s) \leq \frac{\operatorname{Im}(sw(s))}{\sin \beta}. \quad (19)$$

In the last inequality of (19), we used the fact that $\sin(\alpha\beta) \geq \sin \beta > 0$ for all $\alpha \in [\frac{\pi}{2\beta}, 1]$ in order to write $g(s) \leq \int_{\frac{\pi}{2\beta}}^1 r^\alpha \mu(\alpha) d\alpha \leq \frac{\int_{\frac{\pi}{2\beta}}^1 r^\alpha \sin(\alpha\beta) \mu(\alpha) d\alpha}{\sin \beta}$. Therefore, if $\operatorname{Im}(sw(s)) \leq \frac{\lambda \sin \beta}{2}$ then we get from (18)-(19) that $\operatorname{Re}(sw(s) + \lambda) \geq \frac{\lambda}{2}$, which yields (15).

We turn now to proving (16). To this end, we infer from (18)-(19) that

$$\frac{\lambda^\nu (\operatorname{Im}(sw(s)))^{1-\nu}}{|sw(s) + \lambda|} \leq \left(\frac{2}{\sin \beta} \right)^\nu, \quad \nu \in [0, 1]. \quad (20)$$

Indeed, we have already noticed that for $\operatorname{Im}(sw(s)) \leq \frac{\lambda \sin \beta}{2}$ we have $\operatorname{Re}(sw(s) + \lambda) \geq \frac{\lambda}{2}$, whence

$$|sw(s) + \lambda| \geq \sqrt{\left(\frac{\lambda}{2}\right)^2 + (\operatorname{Im}(sw(s)))^2} \geq \left(\frac{\lambda}{2}\right)^\nu (\operatorname{Im}(sw(s)))^{1-\nu}, \quad \nu \in [0, 1].$$

On the other hand, if $\operatorname{Im}(sw(s)) > \frac{\lambda \sin \beta}{2}$ then it holds true that

$$|sw(s) + \lambda| \geq \operatorname{Im}(sw(s)) \geq \left(\frac{\lambda \sin \beta}{2}\right)^\nu (\operatorname{Im}(sw(s)))^{1-\nu}, \quad \nu \in [0, 1].$$

Having established (20), we are now in position to prove (16). To do that, we refer once more to (18)-(19) and examine the two cases $f(s) \geq g(s)$ and $f(s) < g(s)$ separately. We start with $f(s) \geq g(s)$, involving $\operatorname{Re}(sw(s)) \geq 0$ in virtue of (18). Thus we get

$$|sw(s) + \lambda| \geq \sqrt{\lambda^2 + |sw(s)|^2} \geq \lambda^\nu |sw(s)|^{1-\nu}, \quad \nu \in [0, 1],$$

since $\lambda \in (0, +\infty)$, which entails (16). On the other hand, if $f(s) < g(s)$ then it holds true that

$$|\operatorname{Re}(sw(s))| = g(s) - f(s) \leq g(s) \leq \frac{\operatorname{Im}(sw(s))}{\sin \beta},$$

from (18)-(19). As a consequence we have $|sw(s)| \leq \sqrt{1 + \left(\frac{1}{\sin \beta}\right)^2} \operatorname{Im}(sw(s))$, which, combined with (20), yields (16). \square

The second result provides for all $s \in \mathbb{C} \setminus \mathbb{R}_-$ and all $\lambda \in (0, +\infty)$, a suitable lower bound on $|sw(s) + \lambda|$, expressed in terms of $|s|$.

Lemma 2.2. *Let $\mu \in L^\infty(0, 1; \mathbb{R}_+)$ fulfill (6). Then, there exists a constant C , depending only on δ , α_0 and μ , such that we have*

$$|sw(s) + \lambda| \geq C \min(|s|^{\alpha_0 - \delta}, |s|^{\alpha_0}), \quad s \in \mathbb{C} \setminus \mathbb{R}_-, \quad \lambda \in (0, +\infty). \quad (21)$$

Proof. Let us first consider the case where $s = re^{i\beta}$ with $r \in (0, +\infty)$ and $\beta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. Since $\cos(\alpha\beta) \geq 0$ for every $\alpha \in [0, 1]$, then we have

$$|sw(s) + \lambda| \geq \int_0^1 r^\alpha \cos(\alpha\beta) \mu(\alpha) d\alpha + \lambda \geq \int_{\alpha_0 - \delta}^{\alpha_0} r^\alpha \cos(\alpha\beta) \mu(\alpha) d\alpha,$$

in virtue of (6), so we obtain (21) with $C := \frac{\delta \mu(\alpha_0)}{2} \cos\left(\frac{\alpha_0 \pi}{2}\right)$. Similarly, if $|\beta| \in (\frac{\pi}{2}, \pi)$ we infer from (6) that

$$|sw(s) + \lambda| \geq \int_0^1 r^\alpha |\sin(\alpha\beta)| \mu(\alpha) d\alpha \geq \int_{\alpha_0 - \delta}^{\alpha_0} r^\alpha |\sin(\alpha\beta)| \mu(\alpha) d\alpha,$$

which yields (21) with $C := \frac{\delta \mu(\alpha_0)}{2} \min\left(\sin\left(\frac{(\alpha_0 - \delta)\pi}{2}\right), \sin(\alpha_0 \pi)\right)$. \square

Finally, we estimate from above the mapping $s \mapsto |w(s)|$ (resp., $s \mapsto |sw(s)|$) by a suitable continuous monotonically decreasing (resp., increasing) function of $|s|$.

Lemma 2.3. *Assume that $\mu \in L^\infty(0, 1)$ is non-negative. Then, for all $s \in \mathbb{C} \setminus \mathbb{R}_-$, we have*

$$|sw(s)| \leq \|\mu\|_{L^\infty(0,1)} \zeta(|s|) \quad \text{and} \quad |w(s)| \leq \|\mu\|_{L^\infty(0,1)} \vartheta(|s|), \quad (22)$$

where:

(a) *The mapping*

$$\zeta : r \mapsto \begin{cases} \frac{r-1}{\log r} & \text{if } r \in (0, 1) \cup (1, +\infty), \\ 1 & \text{if } r = 1, \end{cases}$$

is continuous and monotonically increasing on $(0, +\infty)$;

(b) The function $\vartheta(r) = \frac{\zeta(r)}{r}$ is continuous and monotonically decreasing on $(0, +\infty)$.

Proof. Since $|sw(s)| \leq \int_0^1 |s|^\alpha \mu(\alpha) d\alpha \leq \|\mu\|_{L^\infty(0,1)} \int_0^1 |s|^\alpha d\alpha$ from the very definition of w , we get (22). Next, for all $r \in (0, 1) \cup (1, +\infty)$, it holds true that $\zeta'(r) = \frac{h(r)}{r(\log r)^2}$ with $h(r) := r \log r - r + 1$. Further, since $h'(r) = \log r$, we have $h(r) \geq h(1) = 0$ for all $r \in (0, 1)$, giving $\zeta'(r) \geq 0$ and hence Statement (a). Finally, Statement (b) follows from the identity $\vartheta(r) = \zeta\left(\frac{1}{r}\right)$ for all $r \in (0, +\infty)$. \square

Armed with the three above lemmas, we may now turn to showing that the IBVP (1) admits a unique solution enjoying a Duhamel representation formula.

2.2 A representation formula

For $\varepsilon \in (0, +\infty)$ and $\theta \in \left(\frac{\pi}{2}, \pi\right)$, we introduce the following contour in \mathbb{C} ,

$$\gamma(\varepsilon, \theta) := \gamma_-(\varepsilon, \theta) \cup \gamma_c(\varepsilon, \theta) \cup \gamma_+(\varepsilon, \theta), \quad (23)$$

where

$$\gamma_\pm(\varepsilon, \theta) := \{s \in \mathbb{C}, \arg s = \pm\theta, |s| \geq \varepsilon\} \text{ and } \gamma_c(\varepsilon, \theta) := \{s \in \mathbb{C}, |\arg s| \leq \theta, |s| = \varepsilon\}. \quad (24)$$

It will prove to be useful for describing the weak solution to the IBVP (1), which is the purpose of the following result.

Proposition 2.1. *Let μ , T , u_0 and F be the same as in Theorem 1.1. Then there exists a unique weak solution*

$$u(t) = S_0(t)u_0 + \int_0^t S_1(t-\tau)F(\tau)d\tau, \quad t \in [0, T], \quad (25)$$

to the IBVP (1), where we have set

$$S_0(t)\psi := \frac{1}{2\pi i} \sum_{n=1}^{+\infty} \left(\int_{\gamma(\varepsilon, \theta)} \frac{w(s)}{sw(s) + \lambda_n} e^{st} ds \right) \langle \psi, \varphi_n \rangle_{L^2(\Omega)} \varphi_n, \quad (26)$$

and

$$S_1(t)\psi := \frac{1}{2\pi i} \sum_{n=1}^{+\infty} \left(\int_{\gamma(\varepsilon, \theta)} \frac{1}{sw(s) + \lambda_n} e^{st} ds \right) \langle \psi, \varphi_n \rangle_{L^2(\Omega)} \varphi_n, \quad (27)$$

for all $\psi \in L^2(\Omega)$, the two above integrals being independent of the choice of $\varepsilon \in (0, +\infty)$ and $\theta \in \left(\frac{\pi}{2}, \pi\right)$. Moreover we have $u \in \mathcal{C}((0, T), L^2(\Omega))$, and even $u \in \mathcal{C}([0, T], L^2(\Omega))$ when $u_0 = 0$.

Proof. The proof is divided into two steps, the first one being concerned with the case of a uniformly vanishing source term F , whereas the second one deals with an identically zero initial state u_0 .

1) *Step 1: $F = 0$.* In order to characterize S_0 we assume that $F = 0$ and put

$$Y(s)u_0 := \sum_{n=1}^{\infty} \frac{w(s)}{s^2(sw(s) + \lambda_n)} \langle u_0, \varphi_n \rangle_{L^2(\Omega)} \varphi_n, \quad s \in \mathbb{C} \setminus \mathbb{R}_-. \quad (28)$$

For all $r \in [2, +\infty)$ and $\eta \in \mathbb{R}$, it is clear from (15) and the estimate $|w(r+i\eta)| \leq \frac{\|\mu\|_{L^\infty(0,1)}}{\log r}$, arising from the second inequality of (22) and Statement (b) in Lemma 2.3, that the inequality

$$|\langle Y(r+i\eta)u_0, \varphi_n \rangle_{L^2(\Omega)}| \leq c \frac{|\langle u_0, \varphi_n \rangle_{L^2(\Omega)}|}{\lambda_n(r^2 + \eta^2)} \leq c \frac{|\langle u_0, \varphi_n \rangle_{L^2(\Omega)}|}{\lambda_n(4 + \eta^2)}, \quad n \in \mathbb{N}, \quad (29)$$

holds with $c := \frac{\|\mu\|_{L^\infty(0,1)}}{\log 2}$. Therefore we have

$$\sup_{r \in [2, +\infty)} \|Y(r + i \cdot)u_0\|_{L^k(\mathbb{R}, L^2(\Omega))} < +\infty, \quad k = 1, 2. \quad (30)$$

From (30) with $k = 1$, it then follows that the function

$$\tilde{y}(t) := \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} e^{ts} Y(s+2)u_0 ds = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{it\eta} Y(2+i\eta)u_0 d\eta,$$

is well defined for each $t \in \mathbb{R}$. Moreover, since the mapping $s \mapsto e^{ts} Y(s+2)u_0$ is holomorphic in $\mathbb{C} \setminus (-\infty, -2]$, we have

$$\tilde{y}(t) = \frac{1}{2\pi i} \int_{\rho-i\infty}^{\rho+i\infty} e^{ts} Y(s+2)u_0 ds, \quad \rho \in (0, +\infty). \quad (31)$$

Indeed, for all $R \in (1, +\infty)$ and all $\rho \in (0, +\infty)$, the Cauchy formula yields

$$\int_{-iR}^{iR} e^{ts} Y(s+2)u_0 ds = \int_{\rho-iR}^{\rho+iR} e^{ts} Y(s+2)u_0 ds - \int_{I_{R,\rho}^+ \cup I_{R,\rho}^-} e^{ts} Y(s+2)u_0 ds, \quad (32)$$

with $I_{R,\rho}^+ := [iR, iR + \rho]$ and $I_{R,\rho}^- := [-iR + \rho, -iR]$, and we infer from (29) that

$$\|Y(r + i\eta)u_0\|_{L^2(\Omega)} \leq \frac{c}{4 + \eta^2} \left(\sum_{n=1}^{\infty} |\langle u_0, \varphi_n \rangle_{L^2(\Omega)}|^2 \right)^{1/2} \leq \frac{c}{4 + \eta^2} \|u_0\|_{L^2(\Omega)},$$

for all $r \in (2, +\infty)$ and $\eta \in \mathbb{R}$, hence

$$\left\| \int_{I_{R,\rho}^\pm} e^{ts} Y(s+2)u_0 ds \right\| \leq \left\| \int_0^\rho e^{t(\tau \pm iR)} Y(\tau + 2 \pm iR)u_0 d\tau \right\| \leq \frac{C\rho e^{\rho t}}{4 + R^2} \|u_0\|_{L^2(\Omega)},$$

so we obtain (31) by sending R to infinity in (32).

Further, with reference to (30) with $k = 1$, we infer from (31) that

$$\|\tilde{y}(t)\|_{L^2(\Omega)} = \frac{1}{2\pi} \left\| \int_{-\infty}^{+\infty} e^{t(\rho+i\eta)} Y(\rho+2+i\eta)u_0 d\eta \right\|_{L^2(\Omega)} \leq \frac{e^{\rho t}}{2\pi} \sup_{r \in [2, +\infty)} \|Y(r + i \cdot)u_0\|_{L^1(\mathbb{R}, L^2(\Omega))},$$

uniformly in $t \in \mathbb{R}$ and $\rho \in (0, +\infty)$. Then, depending on whether $t \in (-\infty, 0)$ or $t \in [0, +\infty)$, we send ρ to either infinity or to zero in the right hand side of the above estimate, and get:

$$\|\tilde{y}\|_{L^\infty(\mathbb{R}, L^2(\Omega))} \leq \frac{1}{2\pi} \sup_{r \in [2, +\infty)} \|Y(r + i \cdot)u_0\|_{L^1(\mathbb{R}, L^2(\Omega))} \quad \text{and} \quad \tilde{y}(t) = 0 \quad \text{for all } t \in (-\infty, 0). \quad (33)$$

As a consequence we have $\tilde{y} \in L^\infty(\mathbb{R}, L^2(\Omega)) \cap \mathcal{S}'(\mathbb{R}_+, L^2(\Omega))$.

Moreover, in light of (30), we infer from [23, Theorem 19.2 and the following remark] and the holomorphicity of $s \mapsto Y(s+2)$ in \mathbb{C}_+ , that $\mathcal{L}[\tilde{y}](s) = Y(s+2)$ for all $s \in \mathbb{C}_+$. Therefore, putting

$$y(t) := e^{2t} \tilde{y}(t) = \frac{e^{2t}}{2\pi i} \int_{-i\infty}^{+i\infty} e^{tp} Y(s+2)u_0 ds = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} e^{st} Y(s)u_0 ds, \quad t \in \mathbb{R},$$

we find that

$$\mathcal{L}[y](s) = \mathcal{L}[\tilde{y}](s-2) = Y(s), \quad s \in \{z \in \mathbb{C}; \operatorname{Re} z \in (2, +\infty)\}. \quad (34)$$

Further, since the mapping $s \mapsto e^{ts}Y(s)a$ is holomorphic in $\mathbb{C} \setminus \mathbb{R}_-$, the Cauchy formula yields

$$y(t) = \frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} e^{ts}Y(s)u_0 ds, \quad t \in \mathbb{R}_+, \quad (35)$$

for all $\theta \in (\frac{\pi}{2}, \pi)$ and all $\varepsilon \in (0, 1)$, the contour $\gamma(\varepsilon, \theta)$ being defined by (23)-(24). Here, we used the fact that

$$\lim_{\eta \rightarrow +\infty} \int_{\eta((\tan \theta)^{-1} \pm i)}^{2 \pm i\eta} e^{ts}Y(s)u_0 ds = 0 \text{ in } L^2(\Omega), \quad t \in \mathbb{R}_+.$$

This can be easily deduced from the following basic estimate, arising from Lemma 2.1, the second inequality of (22) and the second statement of Lemma 2.3,

$$\begin{aligned} \left\| \int_{\eta((\tan \theta)^{-1} \pm i)}^{2 \pm i\eta} e^{ts}Y(s)u_0 ds \right\|_{L^2(\Omega)} &= \left\| \int_{\eta(\tan \theta)^{-1}}^2 e^{t(2 \pm i\eta)}Y(r \pm i\eta)u_0 dr \right\|_{L^2(\Omega)} \\ &\leq C e^{2t} (2 - \eta(\tan \theta)^{-1}) \eta^{-2} \|u_0\|_{L^2(\Omega)}, \quad \eta \in (1, +\infty), \end{aligned}$$

where C is positive constant depending only on θ and μ . We turn now to examining the right hand side of (35). To this purpose, we write for all $t \in \mathbb{R}_+$,

$$y(t) = \sum_{\ell \in \{c, \pm\}} y_\ell(t) \text{ with } y_\ell(t) := \frac{1}{2\pi i} \int_{\gamma_\ell(\varepsilon, \theta)} e^{ts}Y(s)u_0 ds \text{ for } \ell \in \{c, \pm\},$$

and we infer from (15), the second inequality of (22) and Statement (b) in Lemma 2.3, that

$$\|y_\pm(t)\|_{L^2(\Omega)} \leq \frac{\|\mu\|_{L^\infty(0,1)}}{\pi \lambda_1 \sin \theta} \|u_0\|_{L^2(\Omega)} \vartheta(\varepsilon) \int_\varepsilon^{+\infty} r^{-2} e^{r(\cos \theta)t} dr \quad (36)$$

and

$$\|y_c(t)\|_{L^2(\Omega)} \leq \frac{\|\mu\|_{L^\infty(0,1)} (2\theta)^{\frac{1}{2}}}{\pi \lambda_1 \sin \theta} \|u_0\|_{L^2(\Omega)} \varepsilon^{-\frac{3}{2}} \vartheta(\varepsilon) e^{\varepsilon t}. \quad (37)$$

For $t \in [0, e]$, we take $\varepsilon = e^{-1}$ in (36)-(37) and get $\|y_\pm(t)\|_{L^2(\Omega)} + \|y_c(t)\|_{L^2(\Omega)} \leq c(\theta, \mu) \|u_0\|_{L^2(\Omega)}$, where $c(\theta, \mu)$ denotes a generic positive constant depending only on θ and μ . Similarly, for $t \in (e, +\infty)$, we choose $\varepsilon = t^{-1}$, take into account that $\vartheta(t^{-1}) \leq t$, and obtain $\|y_\pm(t)\|_{L^2(\Omega)} + \|y_c(t)\|_{L^2(\Omega)} \leq c(\theta, \mu) \|u_0\|_{L^2(\Omega)} t^{\frac{5}{2}}$. Thus, setting $\langle t \rangle := (1 + t^2)^{\frac{1}{2}}$, we find that

$$\|y(t)\|_{L^2(\Omega)} \leq c(\theta, \mu) \|u_0\|_{L^2(\Omega)} \langle t \rangle^{\frac{5}{2}}, \quad t \in \mathbb{R}_+,$$

which entails $t \mapsto \langle t \rangle^{-\frac{5}{2}} y(t) \in L^\infty(\mathbb{R}_+, L^2(\Omega))$ and hence $y \in \mathcal{S}'(\mathbb{R}_+, L^2(\Omega))$, by (33).

Next, as both functions $s \mapsto \mathcal{L}[y](s)$ and $s \mapsto Y(s)$ are holomorphic in \mathbb{C}_+ , (34) yields $\mathcal{L}[y](s) = Y(s)$ for all $s \in \mathbb{C}_+$, by unique continuation. As a consequence the Laplace transform of $u := \partial_t^2 y \in \mathcal{S}'(\mathbb{R}_+, L^2(\Omega))$ reads

$$\mathcal{L}[u](s) = s^2 Y(s) = \sum_{n=1}^{\infty} \frac{w(s)}{sw(s) + \lambda_n} \langle u_0, \varphi_n \rangle_{L^2(\Omega)} \varphi_n, \quad s \in \mathbb{C}_+,$$

which establishes for each $s \in \mathbb{C}_+$ that $\mathcal{L}[u](s)$ is a solution to (5) with $F = 0$. Moreover, since the operator A is positive and $sw(s) \in [0, +\infty)$ for all $s \in (0, +\infty)$, then $A + sw(s)$ is boundedly invertible in $L^2(\Omega)$. Therefore, $\mathcal{L}[u](s) = (A + sw(s))^{-1} w(s) u_0$ is uniquely defined by (5) and the same is true for u , by injectivity of the Laplace transform.

It remains to prove that u is expressed by (25) with $F = 0$. This can be done by noticing for all $s \in \gamma(\varepsilon, \theta)$ that the function $t \mapsto Y(s)e^{st}u_0$ is twice continuously differentiable in \mathbb{R}_+ and for all $t \in (0, +\infty)$ that $s \mapsto s^k Y(s)e^{st}u_0 \in L^1(\gamma(\varepsilon, \theta))$ with $k = 1, 2$. As a matter of fact, we have

$$\|s^k Y(s)e^{st}u_0\|_{L^2(\Omega)} \leq \frac{2\|\mu\|_{L^\infty(0,1)}}{\lambda_1 \sin \theta} \vartheta(\varepsilon) |s|^{k-2} e^{|s|(\cos \theta)t} \|u_0\|_{L^2(\Omega)}, \quad k = 1, 2, \quad s \in \gamma_\pm(\varepsilon, \theta), \quad t \in \mathbb{R}_+,$$

from the definition (28) of Y , Lemma 2.1 and Statement (b) in Lemma 2.3, whence $r \mapsto r^k Y(re^{\pm i\theta})e^{r(\cos \theta)t} \in L^1(\varepsilon, +\infty)$. Therefore, with reference to (26) we deduce from (35) that $u(t) = S_0(t)u_0$ for all $t \in (0, +\infty)$. We turn now to examining the case where $u_0 = 0$.

2) *Step 2:* $u_0 = 0$. For $s \in \mathbb{C} \setminus \mathbb{R}_-$, we introduce the family of bounded operators in $L^2(\Omega)$,

$$\Phi(s)\psi := \sum_{n=1}^{\infty} \frac{1}{s(sw(s) + \lambda_n)} \langle \psi, \varphi_n \rangle_{L^2(\Omega)} \varphi_n, \quad \psi \in L^2(\Omega), \quad (38)$$

and we recall from (21) that

$$\|\Phi(s)\|_{\mathcal{B}(L^2(\Omega))} \leq C \max(|s|^{-1-\alpha_0}, |s|^{-1-\alpha_0+\delta}), \quad s \in \mathbb{C} \setminus \mathbb{R}_-, \quad (39)$$

for some positive constant C , depending only on μ , in such a way that we have

$$\sup_{r \in [1, +\infty)} \|\Phi(r + i\cdot)\|_{L^k(\mathbb{R}, \mathcal{B}(L^2(\Omega)))} < +\infty, \quad k = 1, 2. \quad (40)$$

Thus, by arguing in the same way as in Step 1, we infer from (39)-(40) that

$$\phi(t) := \frac{1}{2\pi i} \int_{1-i\infty}^{1+i\infty} e^{ts} \Phi(s) ds = \begin{cases} 0 & \text{if } t \in (-\infty, 0), \\ \frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} e^{ts} \Phi(s) ds & \text{if } t \in [0, +\infty), \end{cases} \quad (41)$$

for arbitrary $\varepsilon \in (0, 1)$ and $\theta \in (\frac{\pi}{2}, \pi)$, the contour $\gamma(\varepsilon, \theta)$ being still defined by (23)-(24). Moreover, using (39), we find that

$$t \mapsto \langle t \rangle^{-m_0} \phi(t) \in L^\infty(\mathbb{R}, \mathcal{B}(L^2(\Omega))) \text{ with } m_0 := 1 + \alpha_0. \quad (42)$$

As a consequence we have $\phi \in \mathcal{S}'(\mathbb{R}_+, \mathcal{B}(L^2(\Omega)))$ and

$$\mathcal{L}[\phi\psi](s) = \Phi(s)\psi, \quad s \in \mathbb{C}_+, \quad \psi \in L^2(\Omega). \quad (43)$$

Let us denote by \tilde{F} the function $t \mapsto F(t, \cdot)$ extended by zero in $\mathbb{R} \setminus (0, T)$, and put

$$(\phi * \tilde{F})(t) := \int_0^t \phi(t - \tau) \tilde{F}(\tau) d\tau, \quad t \in \mathbb{R}. \quad (44)$$

Evidently, $\phi * \tilde{F} \in \mathcal{S}'(\mathbb{R}_+, L^2(\Omega))$ when $T \in (0, +\infty)$, whereas for $T = +\infty$, the assumption $t \mapsto \langle t \rangle^{-m} F(t, \cdot) \in L^\infty(\mathbb{R}_+, L^2(\Omega))$ yields

$$\left\| (\phi * \tilde{F})(t) \right\|_{L^2(\Omega)} \leq \left\| \langle t \rangle^{-m_0} \phi \right\|_{L^\infty(\mathbb{R}_+, \mathcal{B}(L^2(\Omega)))} \left\| \langle t \rangle^{-m} F \right\|_{L^\infty(\mathbb{R}_+, L^2(\Omega))} \langle t \rangle^{1+m+m_0}, \quad t \in (0, +\infty), \quad (45)$$

and hence $\phi * \tilde{F} \in \mathcal{S}'(\mathbb{R}_+, L^2(\Omega))$ as well. Further, as we have $\inf\{\epsilon \in (0, +\infty); t \mapsto e^{-\epsilon t} \phi(t) \in L^1(0, +\infty; \mathcal{B}(L^2(\Omega)))\} = 0$ by (42) and $\inf\{\epsilon \in (0, +\infty); t \mapsto e^{-\epsilon t} \tilde{F}(t) \in L^1(0, +\infty; L^2(\Omega))\} = 0$ from the assumption $t \mapsto \langle t \rangle^{-m} F(t, \cdot) \in L^\infty(\mathbb{R}_+, L^2(\Omega))$, then it holds true for all $s \in \mathbb{C}_+$ that $\mathcal{L}[\phi](s) = \int_0^{+\infty} \phi(t) e^{-st} dt$ and $\hat{F}(s) := \mathcal{L}[\tilde{F}](s) = \int_0^{+\infty} \tilde{F}(t) e^{-st} dt$. From this and (45) it then

follows that $\mathcal{L}[\phi * \tilde{F}](s) = \mathcal{L}[\phi](s)\mathcal{L}[\tilde{F}](s)$. As a consequence the function $v := \partial_t(\phi * \tilde{F}) \in \mathcal{S}'(\mathbb{R}_+, L^2(\Omega))$ fulfills

$$\mathcal{L}[v](s) = s\mathcal{L}[\phi * \tilde{F}](s) = \sum_{n=1}^{+\infty} \frac{1}{sw(s) + \lambda_n} \langle \hat{F}(s), \varphi_n \rangle_{L^2(\Omega)} \varphi_n, \quad s \in \mathbb{C}_+,$$

in virtue of (43), showing that $\mathcal{L}[v](s)$ is a solution to (5) for all $s \in (0, +\infty)$.

It remains to show that

$$v(t) = \int_0^t S_1(t - \tau) \tilde{F}(\tau) d\tau, \quad t \in (0, +\infty). \quad (46)$$

To do that, we refer to (41) and (44), apply Fubini's theorem, and obtain

$$(\phi * \tilde{F})(t) = \frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} p(s, t) ds \quad \text{with } p(s, t) := \int_0^t e^{(t-\tau)s} \Phi(s) \tilde{F}(\tau) d\tau, \quad t \in (0, +\infty).$$

Therefore, we have $\partial_t p(s, t) = \Phi(s) \tilde{F}(t) + s \int_0^t e^{(t-\tau)s} \Phi(s) \tilde{F}(\tau) d\tau$ for all $s \in \gamma(\varepsilon, \theta)$ and $t \in (0, +\infty)$, and consequently

$$\begin{aligned} \|\partial_t p(s, t)\|_{L^2(\Omega)} &\leq \|\Phi(s)\|_{\mathcal{B}(L^2(\Omega))} \left(\|\tilde{F}(t)\|_{L^2(\Omega)} + \left| s \int_0^t e^{\tau s} d\tau \right| \|\tilde{F}\|_{L^\infty(0, t+1; L^2(\Omega))} \right) \\ &\leq \left(1 + |s| \int_0^t e^{\tau \operatorname{Re} s} d\tau \right) \|\Phi(s)\|_{\mathcal{B}(L^2(\Omega))} \|\tilde{F}\|_{L^\infty(0, t+1; L^2(\Omega))}. \end{aligned}$$

Putting this with (39), we get that $\|\partial_t p(s, t)\|_{L^2(\Omega)} \leq Cq(s, t) \|\tilde{F}\|_{L^\infty(0, t+1; L^2(\Omega))}$ for some positive constant C depending only on μ , and

$$q(s, t) := \begin{cases} \max(|s|^{-1-\alpha_0+\delta}, |s|^{-1-\alpha_0}) |\cos \theta|^{-1} & \text{if } s \in \gamma_\pm(\varepsilon, \theta), \\ \max(\varepsilon^{-1-\alpha_0+\delta}, \varepsilon^{-1-\alpha_0})(1 + e^{t\varepsilon}) & \text{if } s \in \gamma_c(\varepsilon, \theta). \end{cases}$$

As $s \mapsto q(s, t) \in L^1(\gamma(\varepsilon, \theta))$ for each $t \in (0, +\infty)$, then we have

$$\begin{aligned} v(t) = \partial_t(\phi * \tilde{F})(t) &= \frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} \partial_t p(s, t) ds \\ &= \frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} \int_0^t e^{(t-\tau)s} s \Phi(s) \tilde{F}(\tau) d\tau ds + r(\varepsilon, \theta) \tilde{F}(t), \end{aligned} \quad (47)$$

with $r(\varepsilon, \theta) := \frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} \Phi(s) ds$. The next step is to prove that $r(\varepsilon, \theta) = 0$. For $R \in (1, +\infty)$, put $C_R(\theta) := \{Re^{i\beta}, \beta \in [-\theta, \theta]\}$ and $\gamma_R(\varepsilon, \theta) := \{s \in \gamma(\varepsilon, \theta), |s| \leq R\}$. In light of Lemma 2.2, we have for all $n \in \mathbb{N}$,

$$\int_{\gamma_R(\varepsilon, \theta)} \frac{1}{s(sw(s) + \lambda_n)} ds - \int_{C_R(\theta)} \frac{1}{s(sw(s) + \lambda_n)} ds = 0, \quad (48)$$

by the Cauchy theorem, with $\left| \int_{C_R(\theta)} \frac{1}{s(sw(s) + \lambda_n)} ds \right| \leq 2C\theta R^{-\alpha_0+\delta}$, the constant C being independent of R and n . Thus, sending R to infinity in (48), we get that $\int_{\gamma(\varepsilon, \theta)} \frac{1}{s(sw(s) + \lambda_n)} ds = 0$ for every $n \in \mathbb{N}$, whence $\int_{\gamma(\varepsilon, \theta)} \Phi(s) ds = 0$. From this and (47) we get (46) upon applying the Fubini theorem. Hence $u := v|_Q$ is a weak solution to (5) with $u_0 = 0$, expressed by (25) and (27).

Now, with reference to Step 1 and Step 2, the function represented by (25) is a solution to (1) from the superposition principle. Moreover, it is unique by virtue of Remark 1.1.

It remains to show the last claim of Proposition 2.1. This can be done with the aid of Lemmas 2.2 and 2.3, yielding existence of a constant $C = C(\delta, \alpha_0, \mu) > 0$, such that for any fixed $\theta \in (\frac{\pi}{2}, \pi)$ and any $k = 0, 1$, we have

$$\left| \int_{\gamma_0(1, \theta)} \frac{w(s)^k}{sw(s) + \lambda_n} e^{st} ds \right| \leq C e^t,$$

and

$$\begin{aligned} \left| \int_{\gamma_{\pm}(1, \theta)} \frac{w(s)^k}{sw(s) + \lambda_n} e^{st} ds \right| &\leq C \int_1^{+\infty} r^{-(\alpha_0 - \delta)} e^{rt \cos \theta} dr \\ &\leq C \left(\int_0^{+\infty} \rho^{-\alpha_0 + \delta} e^{\rho \cos \theta} d\rho \right) t^{-1 + \alpha_0 - \delta}, \end{aligned}$$

uniformly in $n \in \mathbb{N}$ and $t \in (0, T)$. Indeed, with reference to (26)-(27), it follows readily from the two above estimates that the right-hand side of (25) lies in $\mathcal{C}((0, T), L^2(\Omega))$, and even in $\mathcal{C}([0, T], L^2(\Omega))$ when $u_0 = 0$. This completes the proof of Proposition 2.1. \square

Theorem 1.1 being a straightforward byproduct of Proposition 2.1, we may now turn to proving Theorems 1.2 and 1.3.

3 Improved regularity: proof of Theorems 1.2 and 1.3

The proof of Theorems 1.2 and 1.3 is by means of suitable time-decay estimates for the operators S_j , $j = 1, 2$, introduced in Proposition 2.1, that are established in the coming section.

3.1 Time-decay estimates

We start with S_0 .

Lemma 3.1. *Let the weight function $\mu \in L^\infty(0, 1)$ be non-negative. Then, for all γ and τ in $(0, 1]$, with $\gamma \leq \tau$, there exists a positive constant C , depending only on θ , μ , λ_1 , τ and γ , such that the estimate*

$$\|A^\tau S_0(t)\psi\|_{L^2(\Omega)} \leq C e^T \|\psi\|_{D(A^\gamma)} t^{\gamma - \tau}$$

holds uniformly in $t \in (0, T]$ and $\psi \in D(A^\gamma)$.

Proof. In light of (26), it is enough to estimate

$$E_n(t) := \frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} \frac{w(s)}{sw(s) + \lambda_n} e^{st} ds, \quad (49)$$

for all $t \in (0, T)$ and $n \in \mathbb{N}$. With reference to (23)-(24), we have

$$E_n(t) = \sum_{j \in \{c, \pm\}} E_{n,j}(t) \text{ where } E_{n,j}(t) := \frac{1}{2\pi i} \int_{\gamma_j(\varepsilon, \theta)} \frac{w(s)}{sw(s) + \lambda_n} e^{st} ds \text{ for } j \in \{c, \pm\}. \quad (50)$$

We shall treat each of the three terms $E_{n,j}(t)$, $j \in \{c, \pm\}$, separately.

We start with $E_{n,c}(t)$. As a preamble, we choose $\varepsilon \in (0, 1)$ so small that

$$\varepsilon < \min(\eta \lambda_1, \zeta^{-1}(\eta \lambda_1)), \quad (51)$$

where $\eta := \frac{1}{2\|\mu\|_{L^\infty(0,1)}}$ and ζ^{-1} denotes the function inverse to ζ on $(0, +\infty)$, whose existence is guaranteed by Statement (a) in Lemma 2.3. As a matter of fact we may take

$\varepsilon := \frac{1}{2} \min(1, \eta\lambda_1, \zeta^{-1}(\eta\lambda_1))$ in such a way that ε is entirely determined by λ_1 and $\|\mu\|_{L^\infty(0,1)}$. In light of (22) and (51), we have $|sw(s) + \lambda_n| \geq \frac{\lambda_n}{2}$ for all $s \in \gamma_c(\varepsilon, \theta)$, and consequently

$$|E_{n,c}(t)| \leq \frac{e^T \lambda_1}{\lambda_n}, \quad t \in [0, T], \quad n \in \mathbb{N}. \quad (52)$$

We turn now to estimating $E_{n,\pm}(t)$. Bearing in mind that $\gamma_\pm(\theta, \varepsilon) = \{re^{\pm i\theta}, r \in (\varepsilon, +\infty)\}$, we decompose $E_{n,\pm}$ into the sum

$$E_{n,\pm}(t) = \sum_{k=1,2} E_{n,\pm,k}(t), \quad (53)$$

where $E_{n,\pm,k}(t) := \frac{1}{2\pi i} \int_{I_{n,k}} \frac{e^{\pm i\theta} w(re^{\pm i\theta})}{re^{\pm i\theta} w(re^{\pm i\theta}) + \lambda_n} e^{rte^{\pm i\theta}} dr$ for $k = 1, 2$, $I_{n,1} := (\varepsilon, \eta\lambda_n)$ and $I_{n,2} := (\eta\lambda_n, +\infty)$. Notice from (51) that $\varepsilon \in (0, \eta\lambda_n)$ and hence that the interval $I_{n,1}$ is non-empty. Next, for all $r \in I_{n,1}$, it is clear from (15) that $|re^{\pm i\theta} w(re^{\pm i\theta}) + \lambda_n| \geq \frac{\lambda_n \sin \theta}{2}$, and from the second inequality of (22) combined with Lemma 2.3, that

$$|w(re^{\pm i\theta})| \leq \|\mu\|_{L^\infty(0,1)} \vartheta(r) \leq \|\mu\|_{L^\infty(0,1)} \vartheta(\varepsilon) \leq \|\mu\|_{L^\infty(0,1)} \frac{\zeta(\varepsilon)}{\varepsilon} \leq \|\mu\|_{L^\infty(0,1)} \frac{\eta\lambda_1}{\varepsilon} \leq \frac{\lambda_1}{2\varepsilon}.$$

As a consequence we have

$$|E_{n,\pm,1}(t)| \leq \frac{1}{2\pi} \int_\varepsilon^{\eta\lambda_n} \frac{|e^{\pm i\theta} w(re^{\pm i\theta})|}{|re^{\pm i\theta} w(re^{\pm i\theta}) + \lambda_n|} e^{-rt|\cos \theta|} dr \leq \frac{\lambda_1}{2\pi\varepsilon\lambda_n \sin \theta} \int_\varepsilon^{\eta\lambda_n} e^{-rt|\cos \theta|} dr$$

whence

$$|E_{n,\pm,1}(t)| \leq \frac{C}{\lambda_n t^{\tau-\gamma}} \int_\varepsilon^{\eta\lambda_n} \frac{dr}{r^{\tau-\gamma}} \leq \frac{C}{\lambda_n^{\tau-\gamma} t^{\tau-\gamma}}. \quad (54)$$

Here and in the remaining part of this proof, C denotes a positive constant depending only on θ , $\|\mu\|_{L^\infty(0,1)}$, λ_1 , τ and γ , that may change from line to line. Further, applying (16) with $\nu = 0$, we get that

$$|E_{n,\pm,2}(t)| \leq \frac{1}{2\pi} \int_{\eta\lambda_n}^{+\infty} \frac{|re^{\pm i\theta} w(re^{\pm i\theta})|}{|re^{\pm i\theta} w(re^{\pm i\theta}) + \lambda_n|} e^{-rt|\cos \theta|} \frac{dr}{r} \leq \frac{1}{\pi \sin \theta} \int_{\eta\lambda_n}^{+\infty} e^{-rt|\cos \theta|} \frac{dr}{r},$$

which entails

$$|E_{n,\pm,2}(t)| \leq \frac{C}{t^{\tau-\gamma}} \int_{\eta\lambda_n}^{+\infty} \frac{dr}{r^{1+\tau-\gamma}} \leq \frac{C}{\lambda_n^{\tau-\gamma} t^{\tau-\gamma}}. \quad (55)$$

Now, putting (50) and (52)–(55) together, we find that

$$\lambda_n^\tau |E_n(t)| \leq C e^T \lambda_n^\gamma t^{\gamma-\tau}, \quad t \in (0, T],$$

Therefore, for all $\psi \in D(A^\gamma)$ and all $t \in (0, T]$, we have

$$\|A^\tau S_0(t)\psi\|_{L^2(\Omega)}^2 = \sum_{n=1}^{\infty} \lambda_n^{2\tau} |E_n(t)|^2 |\langle \psi, \varphi_n \rangle_{L^2(\Omega)}|^2 \leq C^2 e^{2T} t^{2(\gamma-\tau)} \sum_{n=1}^{\infty} \lambda_n^{2\gamma} |\langle \psi, \varphi_n \rangle_{L^2(\Omega)}|^2,$$

which yields the desired result. \square

We turn now to examining the time-evolution of the operator S_1 .

Lemma 3.2. *Assume that $\mu \in L^\infty(0,1)$ is non-negative and satisfies (6). Then, for all $\kappa \in [0,1)$ and all $\beta \in (1 - \alpha_0(1 - \kappa), 1)$, there exists a positive constant C , depending only on θ , λ_1 , μ , κ and β , such that the estimate*

$$\|A^\kappa S_1(t)\psi\|_{L^2(\Omega)} \leq C e^T \|\psi\|_{L^2(\Omega)} t^{-\beta}, \quad (56)$$

holds uniformly in $t \in (0, T]$ and $\psi \in L^2(\Omega)$.

Proof. We proceed as in the proof of Lemma 3.1, the main novelty being that the function $t \in (0, T] \mapsto E_n(t)$, defined by (49) for all $n \in \mathbb{N}$, is replaced by

$$G_n(t) := \frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} \frac{1}{sw(s) + \lambda_n} e^{st} ds. \quad (57)$$

With reference to (23)-(24), we write

$$G_n(t) = \sum_{j \in \{c, \pm\}} G_{n,j}(t) \text{ with } G_{n,j}(t) := \frac{1}{2\pi i} \int_{\gamma_j(\varepsilon, \theta)} \frac{1}{sw(s) + \lambda_n} e^{st} ds \text{ for } j \in \{c, \pm\}, \varepsilon \in (0, 1), \quad (58)$$

and set $\varepsilon = \frac{1}{2} \min(1, \eta\lambda_1, \zeta^{-1}(\eta\lambda_1))$, where $\eta = \frac{1}{2\|\mu\|_{L^\infty(0,1)}}$, as in the proof of Lemma 3.1. Thus, using (22) and (51), we find upon arguing as in the derivation of (52), that

$$|G_{n,c}(t)| \leq \frac{2e^T}{\lambda_n}, \quad t \in (0, T], \quad n \in \mathbb{N}. \quad (59)$$

The rest of the proof is to estimate $G_{n,\pm}(t)$. Firstly, we split $G_{n,\pm}(t)$ into the sum

$$G_{n,\pm}(t) = \sum_{k=1,2} G_{n,\pm,k}(t), \quad (60)$$

where $G_{n,\pm,k}(t) := \frac{1}{2\pi i} \int_{I_{n,k}} \frac{e^{\pm i\theta}}{re^{\pm i\theta} w(re^{\pm i\theta}) + \lambda_n} e^{re^{\pm i\theta} t} dr$ for $k = 1, 2$, $I_{n,1} = (\varepsilon, \eta\lambda_n)$ and $I_{n,2} = (\eta\lambda_n, +\infty)$. By mimicking the proof of (54), we obtain for all $t \in (0, T]$ and all $n \in \mathbb{N}$, that

$$|G_{n,\pm,1}(t)| \leq \frac{1}{\pi\lambda_n \sin\theta} \int_\varepsilon^{\eta\lambda_n} e^{-r|\cos\theta|t} dr \leq \frac{C}{\lambda_n^\kappa t^\kappa}, \quad (61)$$

where C is a positive constant depending only on θ , $\|\mu\|_{L^\infty(0,1)}$ and κ . Next, we estimate $G_{n,\pm,2}(t)$ by applying (16) with $\nu = \kappa$,

$$\begin{aligned} |G_{n,\pm,2}(t)| &\leq \frac{1}{2\pi\lambda_n^\kappa} \int_{\eta\lambda_n}^{+\infty} \frac{\lambda_n^\kappa |re^{\pm i\theta} w(re^{\pm i\theta})|^{1-\kappa}}{|re^{\pm i\theta} w(re^{\pm i\theta}) + \lambda_n|} \frac{e^{-r|\cos\theta|t}}{|re^{\pm i\theta} w(re^{\pm i\theta})|^{1-\kappa}} dr \\ &\leq \frac{1}{\lambda_n^\kappa \sin\theta} \int_{\eta\lambda_n}^{+\infty} \frac{e^{-r|\cos\theta|t}}{|re^{\pm i\theta} w(re^{\pm i\theta})|^{1-\kappa}} dr, \end{aligned} \quad (62)$$

and bounding from below with the aid of (6) the denominator of the integrand in the last integral. Indeed, we have $|re^{\pm i\theta} w(re^{\pm i\theta})| \geq |\operatorname{Im}(re^{\pm i\theta} w(re^{\pm i\theta}))| \geq \int_0^1 r^\alpha \sin(\alpha\theta) \mu(\alpha) d\alpha$ for all $r \in (\eta\lambda_n, +\infty)$, hence $|re^{\pm i\theta} w(re^{\pm i\theta})| \geq \min(\sin((\alpha_0 - \delta)\theta), \sin(\alpha_0\theta)) \frac{\mu(\alpha_0)}{2} \int_{\alpha_0 - \delta}^{\alpha_0} r^\alpha d\alpha$. Consequently there exists a positive constant c_0 , depending only on θ , μ and ρ , such that

$$|re^{\pm i\theta} w(re^{\pm i\theta})| \geq c_0 r^\rho, \quad \rho \in (\alpha_0 - \delta, \alpha_0). \quad (63)$$

Next, applying (63) with $\rho \in \left(\max\left(\alpha_0 - \delta, \frac{1-\beta}{1-\kappa}\right), \alpha_0\right)$ for some fixed $\beta \in (1 - \alpha_0(1 - \kappa), 1)$, we infer from (62) that

$$|G_{n,\pm,2}(t)| \leq \frac{C}{\lambda_n^\kappa t^\beta} \int_{\eta\lambda_n}^{+\infty} \frac{dr}{r^{\beta+\rho(1-\kappa)}} \leq \frac{C}{\lambda_n^\kappa t^\beta}, \quad (64)$$

where C is a positive constant depending only on θ , λ_1 , μ , κ and β . Now, putting (58)–(61) and (64) together, we obtain that

$$|G_n(t)| \leq \frac{Ce^T}{\lambda_n^\kappa t^\beta}, \quad t \in (0, T], \quad n \in \mathbb{N}. \quad (65)$$

Finally, recalling (27), (57) and (65), we end up getting for all $\psi \in L^2(\Omega)$ that

$$\|A^\kappa S_1(t)\psi\|_{L^2(\Omega)}^2 = \sum_{n=1}^{\infty} \lambda_n^{2\kappa} |G_n(t)|^2 |\langle \psi, \varphi_n \rangle_{L^2(\Omega)}|^2 \leq C^2 e^{2T} t^{-2\beta} \sum_{n=1}^{\infty} |\langle \psi, \varphi_n \rangle_{L^2(\Omega)}|^2,$$

and the desired result follows from this and the Parseval identity. \square

Having established Lemmas 3.1 and 3.2, we may now prove Theorem 1.2.

3.2 Proof of Theorem 1.2

We examine the two cases $F = 0$ and $u_0 = 0$ separately.

a) We first assume that $u_0 \in D(A^\gamma)$ for some $\gamma \in (0, 1]$, and that $F = 0$. Then, by applying Lemma 3.1 with $\tau = 1$, we get that

$$\|AS_0(t)u_0\|_{L^2(\Omega)} \leq C e^T \|A^\gamma u_0\|_{L^2(\Omega)} t^{\gamma-1}, \quad t \in (0, T].$$

This and (25) yield (7) in virtue of the equivalence of the norms in $H^2(\Omega)$ and in $D(A)$.

Let us now prove (8). We stick with the notations used in the proof of Lemmas 3.1 and 3.2 and establish for every $\beta \in (1 - \alpha_0\gamma, 1)$ that

$$\left| \frac{dE_n(t)}{dt} \right| \leq C e^T \lambda_n^\gamma t^{-\beta}, \quad t \in (0, T], \quad n \in \mathbb{N}, \quad (66)$$

where C is the constant appearing in (65). This can be done with the aid of (49), (57) and the Cauchy theorem, involving

$$\frac{dE_n(t)}{dt} = \frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} \frac{sw(s)}{sw(s) + \lambda_n} e^{st} ds = -\frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} \frac{\lambda_n}{sw(s) + \lambda_n} e^{st} ds = -\lambda_n G_n(t),$$

upon applying the estimate (65) with $\kappa = 1 - \gamma$. In light of (25)-(26) and (66), we find that $\partial_t u \in \mathcal{C}((0, T), L^2(\Omega))$ fulfills

$$\|\partial_t u(t)\|_{L^2(\Omega)}^2 = \sum_{n=1}^{\infty} \left| \frac{dE_n(t)}{dt} \right|^2 |\langle u_0, \varphi_n \rangle_{L^2(\Omega)}|^2 \leq C^2 e^{2T} t^{-2\beta} \sum_{n=1}^{\infty} \lambda_n^{2\gamma} |\langle u_0, \varphi_n \rangle_{L^2(\Omega)}|^2,$$

which leads to (8).

b) Suppose that $F \in L^1(0, T; L^2(\Omega))$ and $u_0 = 0$. We recall for all $\kappa \in [0, 1)$ and $\beta \in (1 - \alpha_0(1 - \kappa), 1)$ from Lemma 3.2, that the estimate

$$\|A^\kappa S_1(t - \tau)F(\tau)\|_{L^2(\Omega)} \leq C e^T (t - \tau)^{-\beta} \|F(\tau)\|_{L^2(\Omega)}, \quad t \in (0, T], \quad \tau \in [0, t),$$

holds for some constant $C \in (0, +\infty)$, independent of T , t and τ . Therefore we have

$$\|A^\kappa u(t)\|_{L^2(\Omega)} \leq C e^T (\tau^{-\beta} * \|F(\tau)\|_{L^2(\Omega)})(t), \quad t \in (0, T],$$

by (25), where $*$ stands for the convolution operator. An application of Young's inequality then yields

$$\left(\int_0^T \|u(t)\|_{D(A^\kappa)}^p dt \right)^{\frac{1}{p}} \leq C e^T \left(\int_0^T t^{-\beta p} dt \right)^{\frac{1}{p}} \int_0^T \|F(t)\|_{L^2(\Omega)} dt, \quad p \in \left[1, \frac{1}{\beta} \right],$$

and (9) follows from this and the equivalence of the norms in $H^{2\kappa}(\Omega)$ and in $D(A^\kappa)$.

3.3 Proof of Theorem 1.3

The derivation of Theorem 1.3 essentially relies on the following technical result, whose proof is postponed to the appendix.

Lemma 3.3. *Assume that $\mu \in \mathcal{C}([0, 1], \mathbb{R}_+)$ fulfills the condition (6) and let G_n , for $n \in \mathbb{N}$, be defined by (57). Then, we have*

$$G_n(t) = -\frac{1}{\pi} \int_0^{+\infty} \Phi_n(r) e^{-rt} dr, \quad t \in (0, T), \quad (67)$$

with

$$\Phi_n(r) := \frac{\int_0^1 r^\alpha \sin(\pi\alpha) \mu(\alpha) d\alpha}{\left(\int_0^1 r^\alpha \cos(\pi\alpha) \mu(\alpha) d\alpha + \lambda_n \right)^2 + \left(\int_0^1 r^\alpha \sin(\pi\alpha) \mu(\alpha) d\alpha \right)^2}. \quad (68)$$

Moreover, there exists a positive constant C , such that we have

$$\int_0^{+\infty} \frac{\Phi_n(r)}{r} dr \leq \frac{C}{\lambda_n}, \quad n \in \mathbb{N}. \quad (69)$$

As a matter of fact we know from (25) that

$$\|u(t)\|_{D(A)}^2 = \left\| \int_0^t AS_1(t-\tau)F(\tau) d\tau \right\|_{L^2(\Omega)}^2 = \sum_{n=1}^{\infty} \lambda_n^2 \left| \int_0^t G_n(t-\tau) \langle F(\tau), \varphi_n \rangle_{L^2(\Omega)} d\tau \right|^2, \quad t \in (0, T),$$

and for each $n \in \mathbb{N}$ we have

$$\begin{aligned} \int_0^T \left| \int_0^t G_n(t-\tau) \langle F(\tau), \varphi_n \rangle_{L^2(\Omega)} d\tau \right|^2 dt &= \|G_n * \langle F(\cdot), \varphi_n \rangle_{L^2(\Omega)}\|_{L^2(0,T)}^2 \\ &\leq \|G_n\|_{L^1(0,T)}^2 \|\langle F(\cdot), \varphi_n \rangle_{L^2(\Omega)}\|_{L^2(0,T)}^2, \end{aligned}$$

by Young's inequality, whence

$$\|u\|_{L^2(0,T;D(A))}^2 \leq \sum_{n=1}^{\infty} \lambda_n^2 \|G_n\|_{L^1(0,T)}^2 \|\langle F(\cdot), \varphi_n \rangle_{L^2(\Omega)}\|_{L^2(0,T)}^2. \quad (70)$$

Thus, we are left with the task of estimating $\|G_n\|_{L^1(0,T)}$. This can be done by combining Lemma 3.3 with Fubini's theorem, in the same way as in the derivation of [17, Theorem 2.1]. We get for every $n \in \mathbb{N}$ that

$$\|G_n\|_{L^1(0,T)} = \frac{1}{\pi} \int_0^{+\infty} \Phi_n(r) \left(\int_0^T e^{-rt} dt \right) dr \leq \frac{1}{\pi} \int_0^{+\infty} \frac{\Phi_n(r)}{r} dr \leq \frac{C}{\lambda_n},$$

the constant C being independent of n . This and (70) yield

$$\|u\|_{L^2(0,T;D(A))}^2 \leq C \sum_{n=1}^{+\infty} \int_0^T |\langle F(t), \varphi_n \rangle_{L^2(\Omega)}|^2 dt \leq C \|F\|_{L^2(Q)}^2,$$

with the aid of the Parseval identity, which establishes the result.

4 Time analyticity: proof of Theorem 1.4

With reference to Duhamel's representation formula (25) of the weak solution to (1), we may treat the two cases $F = 0$ and $u_0 = 0$ separately. We stick with the notations of Section 2.2 and for any open connected set \mathcal{O} in \mathbb{R} or \mathbb{C} , we denote by $\mathcal{A}(\mathcal{O}, L^2(\Omega))$ the space of $L^2(\Omega)$ -valued analytic functions in \mathcal{O} .

1) *Step 1: $F = 0$.* In light of Proposition 2.1, it is enough to show that the mapping $t \in (0, +\infty) \mapsto S_0(t)u_0$, where S_0 is defined by (26), is extendable to an analytic map of \mathcal{O} into $L^2(\Omega)$, where $\mathcal{O} := \{re^{i\omega}; r \in (0, +\infty), \omega \in (-\theta_1, \theta_1)\}$ for some arbitrary $\theta_1 \in (0, \theta - \frac{\pi}{2}) \cap (0, \pi - \theta)$.

We start by noticing that $|\pm\theta + \omega| \in (\theta - \theta_1, \theta + \theta_1) \subset (\frac{\pi}{2}, \pi)$ for all $\omega \in (-\theta_1, \theta_1)$ and hence that $\cos(\pm\theta + \omega) \leq \cos(\theta - \theta_1)$. Thus we have $|e^{zs}| \leq e^{|z||s|\cos(\theta - \theta_1)}$ for all $z \in \mathcal{O}$ and all $s \in \gamma_{\pm}(\varepsilon, \theta)$, by (24), and (28) then yields $\|e^{zs}s^2Y(s)u_0\|_{L^2(\Omega)} \leq \frac{2\|\mu\|_{L^\infty(0,1)}}{\lambda_1 \sin \theta} \vartheta(\varepsilon) e^{|z||s|\cos(\theta - \theta_1)}$ with the aid of Lemma 2.1 and Statement (b) in Lemma 2.3. Thus, taking into account that $\cos(\theta - \theta_1) \in (-1, 0)$, we get for any compact set $K \subset \mathcal{O}$ that

$$\|e^{zs}s^2Y(s)u_0\|_{L^2(\Omega)} \leq \frac{2\|\mu\|_{L^\infty(0,1)}}{\lambda_1 \sin \theta} \vartheta(\varepsilon) e^{d_K|s|\cos(\theta - \theta_1)}, \quad z \in K, \quad s \in \gamma_{\pm}(\varepsilon, \theta), \quad (71)$$

where $d_K := \min\{|z|, z \in K\} \in (0, +\infty)$. Next, since $z \mapsto e^{zs}s^2Y(s)u_0 \in \mathcal{A}(\mathcal{O}, L^2(\Omega))$ for all $s \in \gamma_{\pm}(\varepsilon, \theta)$, we derive from (71) that $z \mapsto \frac{1}{2\pi i} \int_{\gamma_{\pm}(\varepsilon, \theta)} e^{zs}s^2Y(s)u_0 ds \in \mathcal{A}(\mathcal{O}, L^2(\Omega))$. Further, since the path $\gamma_c(\varepsilon, \theta)$ is finitely extended, it is easy to see that $z \mapsto \frac{1}{2\pi i} \int_{\gamma_c(\varepsilon, \theta)} e^{zs}s^2Y(s)u_0 ds \in \mathcal{A}(\mathcal{O}, L^2(\Omega))$. Therefore, using (23) we find that

$$z \mapsto \frac{1}{2\pi i} \int_{\gamma(\varepsilon, \theta)} e^{zs}s^2Y(s)u_0 ds \in \mathcal{A}(\mathcal{O}, L^2(\Omega)).$$

The desired result follows from this, (26) and (28).

2) *Step 2: $u_0 = 0$.* Fix $t_0 \in (0, +\infty)$. We shall prove existence of $r \in (0, +\infty)$ such that $t \mapsto \int_0^t S_1(\tau)F(t - \tau)d\tau$ is extendable to a holomorphic function of $D(t_0, r)$ into $L^2(\Omega)$. Here, S_1 is the operator defined by (27), and for all $z_0 \in \mathbb{C}$ and all $R \in (0, +\infty)$, we denote by $D(z_0, R) := \{z \in \mathbb{C}; |z - z_0| \in [0, R)\}$ the open disk of \mathbb{C} centered at z_0 with radius R .

First, since F is extendable to a holomorphic function of S_ρ into $L^2(\Omega)$, we may assume without loss of generality that $F \in \mathcal{A}(S_\rho, L^2(\Omega))$. Next we pick $\delta_1 \in (0, \frac{\rho}{4}) \cap (0, t_0)$ in such a way that $z - \tau \in D(0, \rho) \subset S_\rho$ for all $(z, \tau) \in D(t_0, 2\delta_1) \times D(t_0, 2\delta_1)$ and obtain that the mapping

$$\tau \mapsto s\Phi(s)e^{\tau s}F(z - \tau, \cdot) \in \mathcal{A}(D(t_0, 2\delta_1), L^2(\Omega)), \quad z \in D(t_0, 2\delta_1), \quad (72)$$

where Φ the same as in (38).

Further, for $t_1 \in (t_0 - \delta_1, t_0)$ fixed, since $[t_1, z] \subset D(t_0, 2\delta_1)$ for all $z \in D(t_0, 2\delta_1)$, we see from (72) that

$$U_*(s, z) := \int_{[t_1, z]} s\Phi(s)e^{\tau s}F(z - \tau, \cdot)d\tau, \quad s \in \mathbb{C} \setminus \mathbb{R}_-, \quad z \in D(t_0, 2\delta_1) \quad (73)$$

is well defined. Moreover, for all $z \in D(t_0, \delta_1)$ and all $h \in D(0, \delta_1) \setminus \{0\}$, we have

$$\begin{aligned}
& \frac{U_*(s, z+h) - U_*(s, z)}{h} \\
&= \frac{1}{h} \left(\int_{[t_1, z+h]} s\Phi(s)e^{\tau s} F(z+h-\tau, \cdot) d\tau - \int_{[t_1, z]} s\Phi(s)e^{\tau s} F(z+h-\tau, \cdot) d\tau \right) \\
& \quad + \int_{[t_1, z]} s\Phi(s)e^{\tau s} \frac{F(z+h-\tau, \cdot) - F(z-\tau, \cdot)}{h} d\tau \\
&= \int_{[z, z+h]} s\Phi(s)e^{\tau s} F(z+h-\tau, \cdot) d\tau + \int_{[t_1, z]} s\Phi(s)e^{\tau s} \frac{F(z+h-\tau, \cdot) - F(z-\tau, \cdot)}{h} d\tau,
\end{aligned}$$

and hence

$$\lim_{h \rightarrow 0} \frac{U_*(s, z+h) - U_*(s, z)}{h} = s\Phi(s)e^{zs} F(0, \cdot) + \int_{[t_1, z]} s\Phi(s)e^{\tau s} \partial_z F(z-\tau, \cdot) d\tau$$

in $L^2(\Omega)$. As a consequence $z \mapsto U_*(s, z) \in \mathcal{A}(D(t_0, \delta_1), L^2(\Omega))$ for all $s \in \mathbb{C} \setminus \mathbb{R}_-$.

Furthermore, using that $[t_1, z] \subset D(t_0, \delta_1) \subset \mathcal{O}$ for each $z \in D(t_0, \delta_1)$, we derive from (73) that

$$\|U_*(s, z)\|_{L^2(\Omega)} \leq 2\delta_1 |s| \|\Phi(s)\|_{\mathcal{B}(L^2(\Omega))} \|F\|_{L^\infty(D(0, 2\delta_1), L^2(\Omega))} e^{d(t_0, \delta_1)|s| \cos(\theta \pm \theta_1)}, \quad s \in \gamma_\pm(\varepsilon, \theta),$$

with $d(t_0, \delta_1) := \inf\{|z|; z \in D(t_0, \delta_1)\} \in (0, +\infty)$. Thus, bearing in mind that $\theta \pm \theta_1 \in (\frac{\pi}{2}, \pi)$, we infer from this and from (39) that

$$\|U_*(s, z)\|_{L^2(\Omega)} \leq C |s|^{-1-\alpha_0+\delta}, \quad z \in D(t_0, \delta_1), \quad s \in \gamma_\pm(\varepsilon, \theta),$$

for some positive constant C which is independent of s and z . Therefore $z \mapsto \int_{\gamma_\pm(\varepsilon, \theta)} U_*(s, z) ds \in \mathcal{A}(D(t_0, \delta_1), L^2(\Omega))$ and hence

$$z \mapsto \int_{\gamma(\varepsilon, \theta)} U_*(s, z) ds \in \mathcal{A}(D(t_0, \delta_1), L^2(\Omega)). \quad (74)$$

Similarly, by setting

$$U_\#(s, z) := \int_{[0, t_1]} s\Phi(s)e^{\tau s} F(z-\tau) d\tau, \quad s \in \mathbb{C} \setminus \mathbb{R}_-, \quad z \in D(t_0, \delta_1), \quad (75)$$

and arguing as above, we find that

$$z \mapsto \int_{\gamma(\varepsilon, \theta)} U_\#(s, z) ds \in \mathcal{A}(D(t_0, \delta_1), L^2(\Omega)). \quad (76)$$

Finally, putting (25), (27), (38), (73) and (75) together, we obtain that the solution u to (1) reads

$$u(t, \cdot) = \int_0^t S_1(s) F(t-s) ds = \frac{1}{2\pi i} \left(\int_{\gamma(\varepsilon, \theta)} U_*(s, t) ds + \int_{\gamma(\varepsilon, \theta)} U_\#(s, t) ds \right), \quad t \in (t_0 - \delta_1, t_0 + \delta_1),$$

and consequently $u \in \mathcal{A}((t_0 - \delta_1, t_0 + \delta_1), L^2(\Omega))$ by (74) and (76). Since t_0 is arbitrary in $(0, +\infty)$, this entails that $u \in \mathcal{A}((0, +\infty), L^2(\Omega))$, completing the proof of Theorem 1.4.

5 Lipschitz stability: proof of Theorem 1.5

(6) (11)

The strategy of the proof of the stability inequality (12) essentially follows the lines of the derivation of [15, Theorem 2.3] and boils down to the estimates (7)-(8) established in Theorem 1.2.

For the sake of notational simplicity, we start by rewriting the IBVP (1) with $F = 0$ as

$$\begin{cases} \mathbb{D}_t^{(\mu)} u = L_{a,q} u & \text{in } Q, \\ u(0, \cdot) = u_0 & \text{in } \Omega, \\ u = 0 & \text{on } \Sigma, \end{cases} \quad (77)$$

where $L_{a,q}u(x, t) := \operatorname{div}(a(x)\nabla u(x, t)) + q(x)u(x, t)$ is associated with the diffusion matrix a and the electric potential q . Since $u_0 \in D(A)$, we infer from Theorem 1.2 and the energy estimate (7) with $\gamma = 1$ that u and the solution \tilde{u} to (77) where $(\tilde{\mu}, L_{\tilde{a}, \tilde{q}})$ is substituted for $(\mu, L_{a,q})$, belong to $L^\infty(0, T; H^2(\Omega))$. Moreover, as u and \tilde{u} are in $W^{1,1}(0, T; L^2(\Omega))$ by virtue of (8), we get that

$$u, \tilde{u} \in W^{1,1}(0, T; L^2(\Omega)) \cap L^\infty(0, T; H^2(\Omega)). \quad (78)$$

Further, setting $v := u - \tilde{u}$, we deduce from (77) that

$$\begin{cases} \mathbb{D}_t^{(\mu)} v = L_{a,q} v + F & \text{in } Q, \\ v(0, \cdot) = 0 & \text{in } \Omega, \\ v = 0 & \text{on } \Sigma, \end{cases} \quad (79)$$

with

$$F := \mathbb{D}_t^{(\tilde{\mu}-\mu)} \tilde{u} + L_{a-\tilde{a}, q-\tilde{q}} \tilde{u}. \quad (80)$$

In view of (11) and the basic estimate $\Gamma(1-\alpha) \geq 1$ for all $\alpha \in (0, 1)$, we get for all $t \in (0, T)$, upon applying (8) with $\beta = 1 - \alpha_1$, that

$$\begin{aligned} \left\| \mathbb{D}_t^{(\tilde{\mu}-\mu)} \tilde{u}(t) \right\|_{L^2(\Omega)} &\leq \int_0^{\alpha_1} |(\tilde{\mu} - \mu)(\alpha)| \left(\int_0^t (t-s)^{-\alpha} \|\partial_t \tilde{u}(s)\|_{L^2(\Omega)} ds \right) d\alpha \\ &\leq C \left(\int_0^{\alpha_1} \left(\int_0^t (t-s)^{-\alpha} s^{-(1-\alpha_1)} ds \right) d\alpha \right) \|\tilde{\mu} - \mu\|_{L^\infty(0,1)}. \end{aligned}$$

Here and in the remaining part of this proof, C denotes a generic positive constant depending only on $M, \alpha_0, \delta, \alpha_1, \kappa$ and p , which may change from line to line. Next, since

$$\begin{aligned} &\int_0^{\alpha_1} \left(\int_0^t (t-s)^{-\alpha} s^{-(1-\alpha_1)} ds \right) d\alpha \\ &\leq \int_0^{\max(0, t-1)} s^{-(1-\alpha_1)} ds + \int_{\max(0, t-1)}^t (t-s)^{-\alpha_1} s^{-(1-\alpha_1)} ds \\ &\leq \frac{T^{\alpha_1}}{\alpha_1} + \int_0^1 (1-\sigma)^{-\alpha_1} \sigma^{-(1-\alpha_1)} d\sigma, \end{aligned}$$

uniformly in $t \in (0, T)$, we obtain that $\mathbb{D}_t^{(\tilde{\mu}-\mu)} \tilde{u} \in L^\infty(0, T; L^2(\Omega))$. Therefore we have $F \in L^\infty(0, T; L^2(\Omega))$ from (78)-(80), so the solution $u - \tilde{u}$ to (79) belongs $L^p(0, T; H^{2\kappa}(\Omega))$ for all $\kappa \in [0, 1)$ and all $p \in \left[1, \frac{1}{1-\alpha_0(1-\kappa)}\right)$, according to Theorem 9 (b), with the estimate

$$\|u - \tilde{u}\|_{L^p(0, T; H^{2\kappa}(\Omega))} \leq C \|F\|_{L^1(0, T; L^2(\Omega))}. \quad (81)$$

We are thus left with the task of estimating the $L^1(0, T; L^2(\Omega))$ -norm of F . To do that we apply (7)-(8) and get that

$$\|\tilde{u}\|_{L^1(0, T; H^2(\Omega))} \leq Ce^{CT}T \text{ and } \|\partial_t \tilde{u}\|_{L^1(0, T; L^2(\Omega))} \leq Ce^{CT}T^{\alpha_0}. \quad (82)$$

Further, using the fact that (a, q) and (\tilde{a}, \tilde{q}) are in $C^1(\overline{\Omega}) \times C^0(\overline{\Omega})$, we obtain

$$\begin{aligned} \|L_{\tilde{a}-a, \tilde{q}-q} \tilde{u}\|_{L^1(0, T; L^2(\Omega))} &\leq \left(\|a - \tilde{a}\|_{C^1(\overline{\Omega})} + \|q - \tilde{q}\|_{C^0(\overline{\Omega})} \right) \|\tilde{u}\|_{L^1(0, T; H^2(\Omega))} \\ &\leq Ce^{CT} \left(\|a - \tilde{a}\|_{C^1(\overline{\Omega})} + \|q - \tilde{q}\|_{C^0(\overline{\Omega})} \right), \end{aligned} \quad (83)$$

with the aid of (82). On the other hand we have

$$\begin{aligned} \left\| \mathbb{D}_t^{(\tilde{\mu}-\mu)} \tilde{u}(t) \right\|_{L^2(\Omega)} &\leq \|\tilde{\mu} - \mu\|_{L^\infty(0, 1)} \int_0^1 \|\partial_t^\alpha \tilde{u}(t)\|_{L^2(\Omega)} d\alpha \\ &\leq \|\tilde{\mu} - \mu\|_{L^\infty(0, 1)} \int_0^1 \frac{1}{\Gamma(1-\alpha)} \left(\int_0^t \frac{\|\partial_\tau \tilde{u}(\tau)\|_{L^2(\Omega)}}{(t-\tau)^\alpha} d\tau \right) d\alpha, \quad t \in (0, T), \end{aligned}$$

from the very definition of $\mathbb{D}_t^{(\mu)}$, with

$$\int_0^T \int_0^t \frac{\|\partial_\tau \tilde{u}(\tau)\|_{L^2(\Omega)}}{(t-\tau)^\alpha} d\tau dt \leq \|\partial_t \tilde{u}\|_{L^1(0, T; L^2(\Omega))} \int_0^T t^{-\alpha} dt.$$

Therefore we have

$$\begin{aligned} \left\| \mathbb{D}_t^{(\tilde{\mu}-\mu)} \tilde{u} \right\|_{L^1(0, T; L^2(\Omega))} &\leq \|\tilde{\mu} - \mu\|_{L^\infty(0, 1)} \|\partial_t \tilde{u}\|_{L^1(0, T; L^2(\Omega))} \int_0^1 \frac{T^{1-\alpha}}{(1-\alpha)\Gamma(1-\alpha)} d\alpha \\ &\leq Ce^{CT} \|\tilde{\mu} - \mu\|_{L^\infty(0, 1)} T^{\alpha_0} \int_0^1 \frac{T^{1-\alpha}}{\Gamma(2-\alpha)} d\alpha, \end{aligned} \quad (84)$$

by (82) and the identity $\Gamma(2-\alpha) = (1-\alpha)\Gamma(1-\alpha)$. Since the mapping $\alpha \mapsto \Gamma(\alpha)$ is uniformly lower bounded by a positive constant in the interval $[1, 2]$, we deduce from (84) that

$$\left\| \mathbb{D}_t^{(\tilde{\mu}-\mu)} \tilde{u} \right\|_{L^1(0, T; L^2(\Omega))} \leq Ce^{CT} \frac{T^{1+\alpha_0}}{|\log T|} \|\tilde{\mu} - \mu\|_{L^\infty(0, 1)}.$$

Putting this together with (80) and (83), we obtain that

$$\|F\|_{L^1(0, T; L^2(\Omega))} \leq Ce^{CT} \left(\frac{T^{1+\alpha_0}}{|\log T|} \|\mu - \tilde{\mu}\|_{L^\infty(0, 1)} + T \left(\|a - \tilde{a}\|_{C^1(\overline{\Omega})} + \|q - \tilde{q}\|_{C^0(\overline{\Omega})} \right) \right).$$

With reference to (81), this entails for all $\kappa \in [0, 1)$ and all $p \in \left[1, \frac{1}{1-\alpha_0(1-\kappa)}\right)$, that

$$\|u - \tilde{u}\|_{L^p(0, T; H^{2\kappa}(\Omega))} \leq Ce^{CT} \left(\frac{T^{1+\alpha_0}}{|\log T|} \|\mu - \tilde{\mu}\|_{L^\infty(0, 1)} + T \left(\|a - \tilde{a}\|_{C^1(\overline{\Omega})} + \|q - \tilde{q}\|_{C^0(\overline{\Omega})} \right) \right),$$

which yields the desired result.

6 Appendix

6.1 Proof of Lemma 3.3

We stick with the notations used in the proof of Lemma 3.2 and recall from (58) that

$$G_{n,+}(t) + G_{n,-}(t) = -\frac{1}{\pi} \int_\varepsilon^{+\infty} \operatorname{Im} \left(\frac{e^{i\theta+rt} e^{i\theta}}{re^{i\theta} w(re^{i\theta}) + \lambda_n} \right) dr, \quad \varepsilon \in (0, 1).$$

Since the integrand in the above integral reads

$$\operatorname{Im} \left(e^{i\theta+rt e^{i\theta}} \right) \operatorname{Re} \left(\frac{1}{r e^{i\theta} w(r e^{i\theta}) + \lambda_n} \right) + \operatorname{Re} \left(e^{i\theta+rt e^{i\theta}} \right) \operatorname{Im} \left(\frac{1}{r e^{i\theta} w(r e^{i\theta}) + \lambda_n} \right)$$

and that $\lim_{\theta \rightarrow \pi} \operatorname{Im} \left(e^{i\theta+rt e^{i\theta}} \right) = \lim_{\theta \rightarrow \pi} e^{rt \cos \theta} \sin(\theta + rt \sin \theta) = 0$, we derive from (58) that

$$G_n(t) = \frac{1}{2\pi i} \lim_{\theta \rightarrow \pi} \int_{\gamma_c(\varepsilon, \theta)} \frac{1}{s w(s) + \lambda_n} e^{st} ds + \frac{1}{\pi} \int_{\varepsilon}^{+\infty} \Phi_n(r) e^{-rt} dr, \quad (85)$$

where $\Phi_n(r) := \operatorname{Im} \left(\frac{1}{r e^{i\pi} w(r e^{i\pi}) + \lambda_n} \right)$ is expressed by (68). Here, we used the fact that $G_n(t) = \frac{1}{2\pi i} \lim_{\theta \rightarrow \pi} \int_{\gamma_c(\varepsilon, \theta)} \frac{1}{s w(s) + \lambda_n} e^{st} ds$, as the integral in the right hand side of (57) is independent of $\theta \in (\frac{\pi}{2}, \pi)$.

Further, since $\int_{\gamma_c(\varepsilon, \theta)} \frac{e^{st}}{s w(s) + \lambda_n} ds = \int_{-\theta}^{\theta} f_n(\varepsilon, \beta) d\beta$ with $f_n(\varepsilon, \beta) := \frac{i \varepsilon e^{i\beta} e^{\varepsilon t e^{i\beta}}}{\int_0^1 \varepsilon^\alpha e^{i\alpha\beta} \mu(\alpha) d\alpha + \lambda_n}$, and $|f_n(\varepsilon, \beta)| \leq \frac{2\varepsilon e^{\varepsilon t}}{\lambda_n}$ for all $\beta \in [-\pi, \pi]$ provided $\varepsilon \in (0, \min(1, (\eta\lambda_1)^{\frac{1}{\alpha}}))$, then we get

$$\lim_{\varepsilon \rightarrow 0} \left(\lim_{\theta \rightarrow \pi} \int_{\gamma_c(\varepsilon, \theta)} \frac{e^{st}}{s w(s) + \lambda_n} ds \right) = 0,$$

by applying the dominated convergence theorem. Thus, bearing in mind that the right hand side of (85) is independent of $\varepsilon \in (0, +\infty)$ since this is the case for the one of (57), we get (67)-(68) by sending ε to 0 in (85).

We turn now to proving (69). To this purpose, for all $n \in \mathbb{N}$, we introduce $a_n \in (0, +\infty)$ such that

$$\int_0^1 a_n^\alpha \mu(\alpha) d\alpha = \frac{\lambda_n}{2}. \quad (86)$$

Notice that the positive real number a_n is well defined for every $n \in \mathbb{N}$, as the mapping $h : r \mapsto \int_0^1 r^\alpha \mu(\alpha) d\alpha$ is one-to-one from $[0, +\infty)$ onto itself. Moreover, we point out for further use that

$$\lim_{n \rightarrow +\infty} a_n = +\infty. \quad (87)$$

This can be understood from the facts that $a_n = h^{-1}(\frac{\lambda_n}{2})$, where h^{-1} denotes the function inverse to h , that the mapping $r \mapsto h(r)$ is increasing on $[0, +\infty)$, and that $\lim_{r \rightarrow +\infty} h(r) = \lim_{n \rightarrow +\infty} \lambda_n = +\infty$. Next, using (86), we get for all $r \in [0, a_n]$ that $\left| \int_0^1 r^\alpha \cos(\pi\alpha) \mu(\alpha) d\alpha \right| \leq \left| \int_0^1 r^\alpha \mu(\alpha) d\alpha \right| \leq \int_0^1 a_n^\alpha \mu(\alpha) d\alpha \leq \frac{\lambda_n}{2}$, whence $\int_0^1 r^\alpha \cos(\pi\alpha) \mu(\alpha) d\alpha + \lambda_n \geq \frac{\lambda_n}{2}$. Putting this together with (68) and the estimate $\sin u \leq u$, which holds true for all $u \in [0, \pi]$, we find for every $r \in (0, a_n]$ that

$$\frac{\Phi_n(r)}{r} \leq \frac{4}{\lambda_n^2} \int_0^1 r^{\alpha-1} \sin(\pi\alpha) \mu(\alpha) d\alpha \leq \frac{4\pi}{\lambda_n^2} \int_0^1 \alpha r^{\alpha-1} \mu(\alpha) d\alpha \leq \frac{4\pi}{\lambda_n^2} \frac{d}{dr} \left(\int_0^1 r^\alpha \mu(\alpha) d\alpha \right).$$

From this and (86), it then follows that

$$\int_0^{a_n} \frac{\Phi_n(r)}{r} dr \leq \frac{4\pi}{\lambda_n^2} \int_0^1 a_n^\alpha \mu(\alpha) d\alpha \leq \frac{2\pi}{\lambda_n}. \quad (88)$$

The next step of the proof boils down to the fact that there exist two positive constants C_0 and R_0 , both of them depending only on μ , such that we have

$$\int_R^{+\infty} \frac{1}{\int_0^1 r^{\alpha+1} \sin(\pi\alpha) \mu(\alpha) d\alpha} dr \leq \frac{C_0}{\int_0^1 R^\alpha \sin(\pi\alpha) \mu(\alpha) d\alpha}, \quad R \in [R_0, +\infty). \quad (89)$$

The proof of (89), which is quite similar to the derivation of L'Hospital's rule, is presented in Section 6.2 for the convenience of the reader. Prior to applying (89) and upon possibly enlarging R_0 , we notice from (6) that we have

$$\int_0^{\alpha_0-\delta} r^\alpha \mu(\alpha) d\alpha \leq \int_{\alpha_0-\delta}^1 r^\alpha \mu(\alpha) d\alpha, \quad r \in [R_0, +\infty). \quad (90)$$

This follows directly from the two basic estimates $\int_0^{\alpha_0-\delta} r^\alpha \mu(\alpha) d\alpha \leq \|\mu\|_{L^\infty(0,1)}(\alpha_0-\delta)r^{\alpha_0-\delta}$ and $\int_{\alpha_0-\delta}^1 r^\alpha \mu(\alpha) d\alpha \geq \int_{\alpha_0-\frac{\delta}{2}}^{\alpha_0-\delta} r^\alpha \mu(\alpha) d\alpha \geq \frac{\delta\mu(\alpha_0)}{4}r^{\alpha_0-\frac{\delta}{2}}$, which are valid for all $r \in (0, +\infty)$.

Now, in light of (87), we pick $N \in \mathbb{N}$ so large that $a_N \geq R_0$ and we apply (89) with $R = a_n$ for all $n \in \mathbb{N}_N := \{n \in \mathbb{N}, n \geq N\}$. With reference to (68), we obtain that

$$\int_{a_n}^{+\infty} \frac{\Phi_n(r)}{r} dr \leq \int_{a_n}^{+\infty} \frac{1}{\int_0^1 r^{\alpha+1} \sin(\pi\alpha)\mu(\alpha) d\alpha} dr \leq \frac{C_0}{\int_0^1 a_n^\alpha \sin(\pi\alpha)\mu(\alpha) d\alpha}, \quad n \in \mathbb{N}_N.$$

In view of (10), this leads to

$$\int_{a_n}^{+\infty} \frac{\Phi_n(r)}{r} dr \leq \frac{C_0}{\int_{\alpha_0-\delta}^{\alpha_1} a_n^\alpha \sin(\pi\alpha)\mu(\alpha) d\alpha} \leq \frac{C}{\int_{\alpha_0-\delta}^{\alpha_1} a_n^\alpha \mu(\alpha) d\alpha} \leq \frac{C}{\int_{\alpha_0-\delta}^1 a_n^\alpha \mu(\alpha) d\alpha}, \quad n \in \mathbb{N}_N, \quad (91)$$

by setting $C := \frac{C_0}{\min(\sin(\pi(\alpha_0-\delta)), \sin(\pi\alpha_1))} \in (0, +\infty)$. Next, applying (90) with $r = a_n$ and $n \in \mathbb{N}_N$, which is permitted since $a_n \geq a_N \geq R_0$ for all $n \in \mathbb{N}_N$, we get that $\int_{\alpha_0-\delta}^1 a_n^\alpha \mu(\alpha) d\alpha \geq \frac{\int_0^1 a_n^\alpha \mu(\alpha) d\alpha}{2}$. Therefore, we have $\int_{a_n}^{+\infty} \frac{\Phi_n(r)}{r} dr \leq \frac{C}{\lambda_n}$ for all $n \in \mathbb{N}_N$, from (86) and (91). This entails (69) since N is finite.

6.2 Proof of the estimate (89)

For $R \in (0, +\infty)$, put $f(R) := \int_R^{+\infty} \frac{1}{\int_0^1 r^{\alpha+1} \sin(\pi\alpha)\mu(\alpha) d\alpha} dr$ and $g(R) := \frac{1}{\int_0^1 R^\alpha \sin(\pi\alpha)\mu(\alpha) d\alpha}$. Evidently, f and g are two positive functions in $(0, +\infty)$ that vanish at infinity:

$$\lim_{R \rightarrow +\infty} f(R) = \lim_{R \rightarrow +\infty} g(R) = 0, \quad (92)$$

and for all $R \in (0, +\infty)$ and all $R_1 \in (R, +\infty)$, we find by applying Rolle's theorem to the function $r \rightarrow (f(r) - f(R_1))(g(R) - g(R_1)) - (f(R) - f(R_1))(g(r) - g(R_1))$ on the interval $[R, R_1]$, that there exists $\xi \in (R, R_1)$ such that we have

$$\frac{f(R) - f(R_1)}{g(R) - g(R_1)} = \frac{f'(\xi)}{g'(\xi)}. \quad (93)$$

Next, since $\int_0^{\alpha_0-\delta} r^\alpha \sin(\pi\alpha)\mu(\alpha) d\alpha \leq (\alpha_0 - \delta)\|\mu\|_{L^\infty(0,1)}r^{\alpha_0-\delta}$ and

$$\int_{\alpha_0-\delta}^1 r^\alpha \sin(\pi\alpha)\mu(\alpha) d\alpha \geq \int_{\alpha_0-\frac{\delta}{2}}^{\alpha_0} r^\alpha \sin(\pi\alpha)\mu(\alpha) d\alpha \geq cr^{\alpha_0-\frac{\delta}{2}},$$

by (6), where $c := \min(\sin((\alpha_0 - \frac{\delta}{2})\pi), \sin(\alpha_0\pi)) \frac{\delta\mu(\alpha_0)}{4} \in (0, +\infty)$, then there is necessarily $R_0 \in [1, +\infty)$, such that

$$\int_0^{\alpha_0-\delta} r^\alpha \sin(\pi\alpha)\mu(\alpha) d\alpha \leq \int_{\alpha_0-\delta}^1 r^\alpha \sin(\pi\alpha)\mu(\alpha) d\alpha, \quad r \in [R_0, +\infty).$$

From this and the identity $\frac{f'(r)}{g'(r)} = \frac{\int_0^1 r^\alpha \sin(\pi\alpha)\mu(\alpha)d\alpha}{\int_0^1 \alpha r^\alpha \sin(\pi\alpha)\mu(\alpha)d\alpha}$, obtained for all $r \in (0, +\infty)$ by direct calculation, it then follows that

$$\frac{f'(r)}{g'(r)} \leq \frac{2 \int_{\alpha_0-\delta}^1 r^\alpha \sin(\pi\alpha)\mu(\alpha)d\alpha}{(\alpha_0 - \delta) \int_{\alpha_0-\delta}^1 r^\alpha \sin(\pi\alpha)\mu(\alpha)d\alpha} \leq \frac{2}{\alpha_0 - \delta}, \quad r \in [R_0, +\infty).$$

Putting this with (93) we get

$$\frac{f(R) - f(R_1)}{g(R) - g(R_1)} = \frac{f'(\xi)}{g'(\xi)} \leq \frac{2}{\alpha_0 - \delta}, \quad R \in [R_0, +\infty),$$

uniformly in $R_1 \in (R, +\infty)$. Thus, taking $R_1 \rightarrow +\infty$ in the above estimate, we obtain

$$\lim_{R_1 \rightarrow +\infty} \frac{f(R) - f(R_1)}{g(R) - g(R_1)} = \frac{f(R)}{g(R)} \leq \frac{2}{\alpha_0 - \delta}, \quad R \in [R_0, +\infty),$$

in virtue of (92), which is the statement of (89).

Acknowledgment

ZL thanks Grant-in-Aid for Research Activity Start-up 16H06712, JSPS, and National Youth Fund 11801326, NSFC, China. YK and ÉS would like to thank the Department of Mathematical Sciences of The University of Tokyo, where part of this article was written, for its kind hospitality.

References

- [1] Allen M, Caffarelli L, Vasseur A. A Parabolic Problem with a Fractional Time Derivative. *Archive for Rational Mechanics and Analysis*, 2016; 221(2): 603–630.
- [2] Benson D A, et al. Application of a fractional advection-dispersion equation. *Water Resource Research*, 2000; 36: 1403–1412.
- [3] Carcione J M, Sanchez-Sesma F J, Luzón F, et al. Theory and simulation of time-fractional fluid diffusion in porous media. *Journal of Physics A: Mathematical and Theoretical*, 2013; 46(34): 345501.
- [4] Chechkin A V, Gorenflo R, Sokolov I M, et al. Distributed order time fractional diffusion equation. *Fractional Calculus and Applied Analysis*, 2003; 6(3): 259–280.
- [5] Fujishiro K, Kian Y. Determination of time dependent factors of coefficients in fractional diffusion equations. *Mathematical Control and Related Fields*, 2016; 6: 251–269.
- [6] Gorenflo R, Luchko Y, Yamamoto M. Time-fractional diffusion equation in the fractional Sobolev spaces. *Fractional Calculus and Applied Analysis*, 2015; 18(3): 799–820.
- [7] Hatno Y, Hatano Y. Dispersive transport of ions in column experiments: an explanation of long-tailed profiles. *Water Resources Research*, 1998; 34(5): 1027–1033.
- [8] Jiang D, Li Z, Liu Y, Yamamoto M. Weak Unique Continuation Property and a Related Inverse Source Problem for Time-Fractional Diffusion-Advection Equations. *Inverse Problems*, 2017; 33: 055013.

- [9] Kian Y, Oksanen L, Soccorsi E, Yamamoto M. Global uniqueness in an inverse problem for time-fractional diffusion equations. *Journal of Differential Equations*, 2018; 264 (2): 1146–1170.
- [10] Kian Y, Soccorsi E, Yamamoto M. A uniqueness result for time-fractional diffusion equations with space-dependent variable order. to appear in *Annales Henri Poincaré*.
- [11] Kian Y, Yamamoto M. On existence and uniqueness of solutions for semilinear fractional wave equations. *Fractional Calculus and Applied Analysis*, 2017; 20(1): 117–138.
- [12] Kochubei A N. Distributed order calculus and equations of ultraslow diffusion. *Journal of Mathematical Analysis and Applications*, 2008; 340(1): 252–281.
- [13] Kubica A, Ryszewska A. Fractional diffusion equation with the distributed order Caputo derivative. to appear in *J. Integral Equations Applications*.
- [14] Levy M, Berkowitz B. Measurement and analysis of non-Fickian dispersion in heterogeneous porous media. *Journal of contaminant hydrology*, 2003; 64(3): 203–226.
- [15] Li Z, Liu Y, Yamamoto M. Initial-boundary value problems for multi-term time-fractional diffusion equations with positive constant coefficients. *Applied Mathematics and Computation*, 2015; 257: 381–397.
- [16] Li Z, Imanuvilov O Y, Yamamoto M. Uniqueness in inverse boundary value problems for fractional diffusion equations. *Inverse Problems* 32; 2016, 015004.
- [17] Li Z, Luchko Y, Yamamoto M. Analyticity of solutions to a distributed order time-fractional diffusion equation and its application to an inverse problem. *Computers & Mathematics with Applications*, 2016; 73(6): 1041–1052.
- [18] Luchko Y. Boundary value problems for the generalized time-fractional diffusion equation of distributed order. *Fractional Calculus and Applied Analysis*, 2009; 12(4): 409–422.
- [19] Mainardi F, Mura A, Pagnini G, et al. Time-fractional diffusion of distributed order. *Journal of Vibration and Control*, 2008; 14(9-10): 1267–1290.
- [20] Meerschaert M M, Nane E, Vellaisamy P. Distributed-order fractional diffusions on bounded domains. *Journal of Mathematical Analysis and Applications*, 2011; 379(1): 216–228.
- [21] Naber M. Distributed order fractional sub-diffusion. *Fractals*, 2004; 12(1): 23–32.
- [22] Nochetto R, Otárola E, Salgado A A PDE Approach to Space-Time Fractional Parabolic Problems *SIAM J. Numer. Anal.*, 54(2), 848–873.
- [23] Rudin W. *Real and complex analysis*. Tata McGraw-Hill Education, 1987.
- [24] Rundell W, Zhang Z. Fractional diffusion: recovering the distributed fractional derivative from overposed data. *Inverse Problems*, 2017; 33: 035008.
- [25] Sakamoto K, Yamamoto M. Initial value /boundary value problems for fractional diffusion-wave equations and applications to some inverse problems. *J. Math. Anal. Appl.*, 2011; 382: 426–447.
- [26] Sokolov I M, Chechkin A V, Klafter J. Distributed-Order Fractional Kinetics. *Acta Physica Polonica B*, 2004; 35: 1323.

- [27] Sun H G, Chen W, Chen Y Q. Variable-order fractional differential operators in anomalous diffusion modeling. *Physica A: Statistical Mechanics and its Applications*, 2009; 388(21): 4586–4592.
- [28] Zacher R. Weak solutions of abstract evolutionary integro-differential equations in Hilbert spaces. *Funkcialaj Ekvacioj*, 2009; 52(1): 1–18.